

インターナショナルセッション（口頭発表） | セッション記号 A（大気海洋・環境科学） | A-CG 大気海洋・環境科学複合領域・一般

[A-CG05_30AM1]Continental-Oceanic Mutual Interaction: Global-scale Material Circulation through River Runoff

コンビーナ:*山敷 庸亮(京都大学大学院総合生存学館)、Behera Swadhin(Climate Variation Predictability and Applicability Research Program Research Institute for Global Change/JAMSTEC, 3173-25 Showa-machi, Yokohama 236-0001)、升本 順夫(海洋研究開発機構)、宮澤 泰正(独立行政法人海洋研究開発機構)、山形 俊男(海洋研究開発機構 アプリケーションラボ)、寶 馨(京都大学防災研究所)、座長:升本 順夫(海洋研究開発機構)、Behera Swadhin(Climate Variation Predictability and Applicability Research Program Research Institute for Global Change/JAMSTEC, 3173-25 Showa-machi, Yokohama 236-0001)、山形 俊男(海洋研究開発機構 アプリケーションラボ)

2014年4月30日(水) 09:00 ~ 10:45 211 (2F)

The main purpose of this session is to promote discussion on mutual interaction between Continental zone and Oceanic zone. The global-scale material circulation induced by River runoff through oceanic general circulation as major topic on Continental-Oceanic Interaction, where the ENSO / IOD influence into continental climate as major topics on Oceanic-Continental Interaction. Numerical simulation and field observation of radionuclide transport from continental zone into ocean and its potential impact is also important topics of this session.

10:35 ~ 10:45

[ACG05-P01_PG]Land-Ocean Mutual Interaction: Sediment Transportation Processes in Coastal Zone Induced by Abukuma River Mouth Runoff

ポスター講演3分口頭発表枠

*TROSELJ Josko¹、YAMASHIKI Yosuke²、TAKARA Kaoru³ (1.Graduate School of Engineering, Kyoto University、2.Graduate School of Advanced Integrated Studies in Human Survivability, Kyoto University、3.Disaster Prevention Research Institute, Kyoto University)

キーワード : River-ocean interaction, Integrated atmosphere-land-ocean model, Sediment transportation of radionuclides, Lagrangian particle tracking model, Near-shore bathymetry influence

1. Introduction

Modeling of a contact zone between a river coming from potentially contaminated basin and an ocean is especially sensitive case for Land-Ocean coupling interaction due to significant risk of major environmental disaster which can occur in the case of contamination of the coastal zone. Therefore, it is of great importance to study and develop integrated modeling approach to comprehend the complex interaction processes in the contact zone in order to minimize disaster risk potential, which can consequently cause undesirable social and economical costs.

2. Objectives

The focus of this study is to promote relevant numerical simulation on Land-Ocean coupling modeling approach applicable for the bay and estuary zone affected by river inflow and associated sediment transportation from the Abukuma river basin in Japan. By conducting several field observations, we found interesting and unusual temporal and spatial distribution of radionuclides within the coastal zone near the river mouth. Sediment transportation processes which have led to the distribution as well as influence of near-shore bathymetry to sediment dispersion are of close interest for the study.

Calculation was conducted by simulating stages and conditions for mechanisms of sediment transport in the coastal zone, from initial deposition onto river bed to final deposition onto ocean floor, with special focus on processes which are occurring during high water periods.

3. Model Description

The modeling approach has been studied by combining river mouth runoff boundary conditions by using Lagrangian particle tracking model for simulating sediment transportation, with coupled atmosphere-ocean-land model (MSSG model, JAMSTEC) which used fine resolution grid, and associated downscaling techniques for oceanic boundary conditions (JCOPE2 model, JAMSTEC) which used coarse resolution grid. We have simulated two different cases, at first circulation of ocean itself in non-equilibrium quasi stationary state, where its dynamics was induced only by its own temperature and salinity data differences among adjacent cells, and at second response of the ocean circulation to inflow from the river outlet, simulated under various boundary conditions and external effects.

ETOPO1, 1 Arc Minute Global Relief Model was chosen as initial database for bathymetry data, while World Ocean Atlas 2005 database was chosen as initial 3D database for temperature, salinity, pressure, and velocity field data. Incompressible Navier-Stokes equation and Yin-Yang grid were used in the calculation of the flow field.

4. Conclusions and follow up

The study is continuous part of the PhD study of the first author, so the results are about to be improved as the course will continue. So far, results neither confirmed nor denied the hypothesis that near-shore bathymetry may have important role in spatial dispersion of radionuclides, so the question still remains open and subject for discussion. Our assumption is that using fine resolution grid within the contact zone between two different fluids should give us better insight into the problem, while simultaneously proper downscaling of outer oceanic boundary conditions and proper coupling with sediment transportation model are needed to be done in order to maintain satisfactory level of simulated physics of processes during the calculation. In follow up of the study, we will try to simulate hydrograph based water wave rather than constant inflow from the river mouth, as well as try to put ocean side into initial dynamic state rather than non-equilibrium quasi stationary state. Also, more focus will be aimed to the physical processes behind mechanisms of radionuclide transportation from the basin towards the river mouth.

Acknowledgements

Dr. Shinichiro Kida, executive supervisor, JAMSTEC

Dr. Keiko Takahashi, MSSG program director, JAMSTEC

Monbukagakusho (MEXT) scholarship, GCOE-HSE program