

---

Oral | Symbol H (Human Geosciences) | H-GM Geomorphology

## [H-GM22\_30AM2]Geomorphology

Convener:\*Hiroshi Shimazu(Department of Geography, Faculty of Geo-Environmental Science, Rissho University), Chiaki T. Oguchi(Geosphere Research Institute, Saitama University), Masayuki Seto(Fukushima Future Center for Regional Revitalization, Fukushima University), Chair:Masayuki Seto(Fukushima Future Center for Regional Revitalization, Fukushima University)

Wed. Apr 30, 2014 11:00 AM - 12:45 PM 422 (4F)

The main subject of this session is interdisciplinary discussion on the whole range of themes relating to geomorphology, especially geomorphic processes, landform development, geomorphological hazards and their mitigation, and relationships among geomorphic processes, other natural phenomena and human activities. All topics on geomorphology with new findings and ideas are welcome. All presentations and discussion of this session are made in Japanese.

---

12:15 PM - 12:30 PM

## [HGM22-P06\_PG]The characteristic of roots distribution on the slopes in Izu-Oshima where landslides were occurred by Typhoon No.26

3-min talk in an oral session

\*Wataru MURAKAMI<sup>1</sup>, Akiho OGAWA<sup>1</sup>, Yasuhiro OGAWA<sup>1</sup>, Hiromu DAIMRU<sup>1</sup> (1.FFPRI)

Keywords:Izu-Oshima, Typhoon No.26, landslide, roots distribution

Large-scale landslides were occurred in Izu-Oshima by Typhoon No.26 on October 16, 2013. By the urgent investigation after the disaster, it is reported that the landslides were occurred in the part within about 1m from the slope surface and the few rhizomes were on the slip surface. We surveyed the distribution of the fallen trees (species, height, the root depth, and the extensions (widths) of the roots) on the slope near the landslide. The surveyed fallen trees were a lot of *Eurya japonica*, and were the order of *Ilex crenata var.hachijoensis*, *Prunus lannesiana var. speciosa*, and *Camellia japonica*. Most of the surveyed fallen trees were about 5-7m in height, and the high one was 10m or less. The root depth of most fallen trees was 60-80cm; however, the root depth had the difference by the tree species. The *Camellia japonica* and the *Prunus lannesiana var. speciosa* had comparatively deep roots. On the other hand, the *Eurya japonica* and the *Ilex crenata var.hachijoensis* tended to be distributed shallowly the root systems. As a factor to which the root systems are not deeply distributed, a peculiar properties of soil situation of the volcano is pointed out. In this survey, the difference of characteristics of the tree species on the surveyed slopes was guessed as a cause, too.