## Oral | Symbol M (Multidisciplinary and Interdisciplinary) | M-AG Applied Geosciences

## [M-AG38\_2PM2]Dynamics of radionuclides emitted from Fukuchima Dai-ichi Nuclear Power Plant in the environment

Convener:\*Kazuyuki Kita(Faculty of Science, Ibaraki University), Yuichi Onda(Center for Research on Isotopes and Environmental Dynamics, University of Tsukuba), Teruyuki Nakajima(Atmosphere and Ocean Research Institute), Yasuhito Igarashi(Atmospheric Environment and Applied Meteorology Research Department, Meteorological Research Institute), Jun Matsumoto(Deaprtment of Geography, Tokyo Metropolitan University), Masatoshi Yamada(Institute of Radiation Emergency Medicine, Hirosaki University), Chisato Takenaka(Graduate School of Bioagricultural Sciences, Nagoya University), masayoshi yamamoto(Low Level Radioactivity Laboratory, Kanazawa University), jota kanda(Tokyo University of Marine Science and Technology), atsushi shinohara(Osaka university), Chair:Motoyoshi Ikeda(Hokkaido University)

Fri. May 2, 2014 4:15 PM - 5:00 PM 501 (5F)

The Great East Japan Earthquake caused the severe accident in TEPCO Fukushima dai-ichi nuclear power plant (FDNPP), leading to emission of huge amount of radionuclides to the environment. They have been transported and diffused by atmospheric motion, depositing them to soil and vegetation. Deposited radionuclides are dynamically shifted in the earth environment; atmosphere, soil, inland water, ocean, and ecosystem. To understand this dynamic shift in the environment and for the long-term prediction of the disaster by the radionuclides, investigation and discussion based on not only the earth sciences including ecology but also on the radiochemistry and other related sciences. In this session, various efforts to understand the dynamic behavior of radionuclides emitted from FDNPP accident in the earth system as well as to predict their influences on the environment. It is expected that this session will offer a good opportunity to discuss radionuclides in the earth environment from wide aspect and to exchange information in various research fields.

## 4:45 PM - 5:00 PM [MAG38-P06\_PG]Secular distribution of radioactive strontium concentration in the atmosphere after after the accident of FD-NPP

3-min talk in an oral session

\*Zi jian ZHANG<sup>1</sup>, Kazuhiko NINOMIYA<sup>1</sup>, Naruhito TAKAHASHI<sup>1</sup>, Yoshiaki YAMAGUCHI<sup>2</sup>, Takashi YOSHIMURA<sup>2</sup>, Takashi SAITO<sup>3</sup>, Kazuyuki KITA<sup>4</sup>, Haruo TSURUTA<sup>5</sup>, Akira WATANABE<sup>6</sup>, Atsushi SHINOHARA <sup>1</sup> (1.Graduate School of Science, Osaka University, 2.Radioisotope Research Center, Osaka University, 3.Faculty of Comprehensive Human Sciences, Shokei Gakuin University, 4.College of Science, Ibaraki University, 5.Atmosphere and Ocean Research Institute, the University of Tokyo, 6.Fukushima University)

1.IntroductionOn March 12, 2011, a large amount of radioactive nuclides have been released into the environment by the nuclear accident at the Fukushima Daiichi Nuclear Power Station. Measurement about radioactive nuclides will give us much information about the accident circumstance. Furthermore, radioactivities in the air dust are critical for estimation of internal exposure. There are many measurements results of I-131, Cs-134, Cs-137 in environment samples. However, in other nuclides, such as the pure beta emitter nuclide Sr-90 has not been measured sufficiently. Sr-90 is considered one of the harmful radioactive nuclides. Therefore, measurement of Sr-90 in the air dust is important for calculating exposure. We developed a new simple and quick strontium isolation technique using solid-

phase extraction for determination Sr-90 in the air dust by liquid scintillation counter (LSC). 2. MethodIn this study, we used 3M EmporeTM Strontium Rad Disk to extract strontium ion from air dust samples. This filter can collect Sr2+ ion efficiently. However, it is known that this filer also catches Pb2+. Natural radioactive nuclide Pb-210 seriously will be interferences in Sr identification in beta ray counting. In this study, cation exchange with EDTA adopted for Sr isolation. We made test experiments with radioactive Sr tracer and obtained that the chemical yield was about 90 %. The time for chemical operation was about 3-4 hours. To determine Sr-90, Cherenkov radiation of Y-90 has been measured by LSC, 1220 QUANTULUSTM Ultra Low Level Liquid Scintillation Spectrometer. With Sr-90 standard solution, we obtained that the Y-90 Cherenkov light detection efficiency was 68.7% and the Sr-90 detection limit was 0.004 Bq. With sequential measurement, the growth curve of Y-90 was described to determinate activity of Sr-90.3.Results We measured Sr-90 in the air dust samples of Fukushima, Hitachi, Kawasaki and Osaka. We chose some air dust samples that have high Cs-137 activity for Sr-90 measurement. Strontium isolation with solid phase extraction was performed. In Hitachi, the Sr-90 activity concentration in air is decreased with time and the ratio of Sr-90/Cs-137 is about 10-3. It is possible that after April, Sr-90 has been the same behavior of Cs-137. We observed a long time variation of Sr-90 air concentration in Hitachi and Fukushima and found that the Sr-90/Cs-137 activity ratio increased over time. We are going to discuss about behaviors of the Sr-90 and Cs-137 in the atmosphere.