Japan Geoscience Union Meeting 2014

(28 April - 02 May 2014 at Pacifico YOKOHAMA, Kanagawa, Japan)

©2014. Japan Geoscience Union. All Rights Reserved.



PEM05-04 Room:424

Time:May 2 10:00-10:15

Long-term observations of MLT zonal wind variations in relation to stratospheric zonal winds over low-latitudes

GRANDHI, Kishore kumar^{1*}; KARANAM, Kishore kumar²; SINGER, Werner¹; ZULICKE, Christoph¹; S, Gurubaran³; GERD, Baumgarten¹; RAMKUMAR, Geetha²; S, Sathishkumar⁴; RAPP, Markus⁵

¹Leibniz Institute of Atmospheric Physics, University of Rostock, Kuhlungsborn, Germany, ²Space Physics Laboratory, Vikram Sarabhai Space Center, Trivandrum, India, ³Indian Institute of Geomagnetism, Navi Mumbai, India, ⁴Equatorial Geophysical Research Laboratory, Indian Institute of Geomagnetism, Tirunelveli, India, ⁵German Aerospace Center Institute of Atmospheric Physics (IPA), Oberpfaffenhofen, Wessling, Germany

Long-term observations from medium-frequency and meteor radars (1993-2012) and rocket soundings (1979-1990 and 2002-2007) are used to study mesosphere lower thermosphere (MLT) zonal wind variations in relation to the stratospheric winds over Northern low-latitudes. The combined dataset provide a complete height profile of amplitude of semiannual oscillation (SAO) up to 100 km, with an exception around 75-80 km. The SAO signal has maxima around 50 km and 82 km and a minimum around 65 km. The MLT zonal winds show remarkable inter-annual variability during spring equinox and much less during fall equinox. Zonal wind mesospheric spring equinox enhancements (MSEE) appear with a periodicity of 2-3 years suggesting a modulation by the quasi-biennial oscillation, which we identified with the strength of stratospheric westward winds. Out of 20 years of observations, the stratospheric westward winds are strong during 11 years (non-MSEE) and weak during 9 years. Six of these years show large MLT winds (MSEE) and 3 years (1999, 2004 and 2006) show small MLT winds (missing-MSEEs). These unexpected small winds occur in years with global circulation anomalies as identified with strong sudden stratospheric warmings and an early spring transition of zonal winds, along with a minor enhancement in the tidal amplitudes.

Keywords: MLT winds, MSAO, Meteor radar, MF radar, QBO