

インターナショナルセッション（口頭発表） | セッション記号 P（宇宙惑星科学） | P-EM 太陽地球系科学・宇宙電磁気学・宇宙環境

## [P-EM06\_30PM2] Study of coupling processes in Sun-Earth system with large radars and large-area observations

コンビーナ:\*山本 衛(京都大学生存圏研究所)、小川 泰信(国立極地研究所)、野澤 悟徳(名古屋大学太陽地球環境研究所)、橋口 浩之(京都大学生存圏研究所)、座長:橋口 浩之(京都大学生存圏研究所)

2014年4月30日(水) 16:15 ~ 18:00 312 (3F)

The Earth accepts huge input of energy and material from the Sun. The Earth's environment is maintained by the balance between their inputs and outputs. It is important to study energy and material transport of the Earth. This is an international session that discusses studies of the coupling processes in the Sun-Earth system based on the projects of large radars and large-area observation network. The facilities and networks included are the Equatorial MU Radar (EMU) in Indonesia to study the whole equatorial atmosphere, the EISCAT\_3D radar system to study detailed structures and elementary processes of the magnetosphere-ionosphere in the polar region, and global observation networks of magnetometers and radio and optical instruments to study the coupling processes with the global scale. We will show outline of the project and discuss sciences by soliciting variety papers. This session is open to the world, and we strongly encourage submission of papers related to other facilities and projects, i.e., atmospheric or incoherent-scatter radars, observation networks, satellites, and simulation or theoretical studies, etc.

17:15 ~ 17:30

## [PEM06-P10\_PG] Naライダー/EISCATレーダー連携観測によって検出した オーロラ起因の Na層変動

ポスター講演3分口頭発表枠

\*津田 卓雄<sup>1</sup>、野澤 悟徳<sup>2</sup>、川原 琢也<sup>3</sup>、川端 哲也<sup>2</sup>、斎藤 徳人<sup>4</sup>、和田 智之<sup>4</sup>、小川 泰信<sup>1</sup>、大山 伸一郎<sup>2</sup>、Hall Chris<sup>5</sup>、堤 雅基<sup>1</sup>、江尻 省<sup>1</sup>、鈴木 臣<sup>2</sup>、高橋 透<sup>2</sup>、中村 卓司<sup>1</sup> (1.国立極地研究所、2.名古屋大学太陽地球環境研究所、3.信州大学工学部、4.理化学研究所、5.トロムソ大学)

キーワード：Naライダー, EISCATレーダー, Na層, オーロラ降下粒子, 電離圏電場

Sodium atom layer is generally distributed at 80-100 km. One of mysterious subjects on high-latitude sodium layers is relationship between auroral particle precipitation and sodium atom layer variation. A previous study suggested a sodium column density decrease during a geomagnetic active period due to that the particle precipitation accompanied by electron density enhancement could induce ionization of sodium atom through their ion-molecule chemistry. Another study pointed a possibility of sodium density increase. For this reason, it is suggested that auroral precipitating particle bombardment on meteoric smoke particles can sputter sodium atoms from the smoke particles. On the other hand, ionospheric electric field, which may become more significant near auroral precipitating regions, could induce ion motions (i.e. can generate sodium ion convergence and/or divergence), and then also could affect generation and/or loss processes of sodium atoms through their ion-molecule chemistry. Thus, for the examination of the causality, it is vitally important to distinguish the effects of auroral particle precipitation and ionospheric electric field. Using a sodium lidar (which was installed in early 2010) and European incoherent scatter (EISCAT) radar at Tromsø, Norway (69.6N, 19.2E), we have investigated, for the first time, that the actual effect of the particle precipitation to the sodium density variations without electric field injection. In the nighttime observation on 24-25 January 2012, we detected a significant decrease of sodium atom density coincided with electron density enhancements (implying strong particle precipitations) and low ion temperatures (implying no electric field injections). These

results strongly suggested that auroral particle precipitations induced sodium atom density decrease. Furthermore we discuss observed time response in the sodium density decrease.