
Oral | Symbol S (Solid Earth Sciences) | S-EM Earth's Electromagnetism

[S-EM36_30PM2]Electrical conductivity, Tectono-electromagnetism

Convener:*Ken'ichi Yamazaki(Disaster Prevention Research Institute, Kyoto University), Noriko Tada(Japan Agency for Marine-Earth Science and Technology), Chair:Noriko Tada(Japan Agency for Marine-Earth Science and Technology), Ken'ichi Yamazaki(Disaster Prevention Research Institute, Kyoto University)

Wed. Apr 30, 2014 4:15 PM - 5:45 PM 413 (4F)

This session is for wide variety of studies on electromagnetism of solid Earth. The topics include electromagnetic phenomena associated with earthquakes and volcanism, electrical conductivity structure, laboratory experiments, results of simulations, new equipments for observation, and methods of data analysis.

5:30 PM - 5:45 PM

[SEM36-P06_PG]Electrical resistivity features of the back-arc areas in the NE Japan subduction zone

3-min talk in an oral session

*Hiroshi ICHIHARA¹, Noriko TADA¹, Kiyoshi BABA², Takafumi KASAYA¹, Masahiro ICHIKI³, Toshiki KAIDA³, Yasuo OGAWA⁴ (1.JAMSTEC, 2.ERI, University of Tokyo, 3.Tohoku University, 4.Tokyo Institute of Technology)

Keywords:back arc, NE Japan subduction zone, magnetotelluric, OBEM, phase tensor

Electrical resistivity in the crust and upper mantle depends on the pore-fluid distribution, salinity, and connectivity of fluid-filled rock pores. Thus imaging of resistivity distribution based on magnetotelluric surveys gives us fundamental information about fluid distribution of subduction zones. Marine magnetotelluric survey is important to understand dynamics of the NE Japan subduction zone because dehydration of subducting Pacific plate occurs under the Japan Sea. In this study, we discuss resistivity distribution around back-arc areas in the NE Japan subduction zone based on the marine MT data. We collected natural EM signals with ocean bottom electro-magnetometers (OBEMs) in the eastern Japan sea area between April and August 2013 by MR13-02A and NT13-18 JAMSTEC scientific cruises. In addition, 3 land MT stations were settled in islands in the Japan Sea (Tobishima, Awashima and Sado islands) between April and October 2013. These recorded time-series data were converted to a frequency-domain impedance tensor based on the BIRRP program [1]. The remote reference technique [2] was applied in the data processing using horizontal magnetic field data from Kakioka Station in the period range between 10 and 20000 seconds. As results, high-quality MT responses and geomagnetic tippers in both the trench and back-arc areas. We calculated phase tensors [3] based on MT impedances by this and previous studies [4] to discuss re-sistivity distribution beneath the back-arc area. The phase tensor ellipse indicates high Φ_{\max} (>65 degrees) and Φ_{\min} (>50 degrees) in the long periods (>8000 seconds). Large β of phase tensor and large amplitude of geomagnetic transfer function are also shown. These features cannot be explained with bathymetry and sediment effects based on the 3-D forward modeling [5]. Thus strong three-dimensionality and deep conductor possibly distributed beneath the Japan sea. In order to discuss detailed resistivity structure, 3-D inversion approaches are required by using a newly developed 3-D MT inversion code for marine data to treat complicated ocean bottom and land topography [6]. **References:** [1] Chave, A. D. and D. J. Thomson, *Geophys. J. Int.* 157, 988-1006 (2004); [2] Gamble, T. D. et al., *Geophysics*, 44, 53-68 (1979); [3] Caldwell, T. G et al., *Geophys. J. Int.* 158, 457-469 (2004); [4] Toh, H. et al., *Geophys Res Lett*, 33, L22309 (2006); [5] Baba, N. et al.,

Geophys. J. Int 158, 392-402 (2002); [6] Tada, N. et al., Earth Planets Space, 64, 10051021 (2012).