Japan Geoscience Union Meeting 2015

(May 24th - 28th at Makuhari, Chiba, Japan)

©2015. Japan Geoscience Union. All Rights Reserved.

PEM11-11

Room:A01



Time:May 25 10:00-10:15

## Airglow image of mesospheric mesoscale wave captured from the International Space Station

HOZUMI, Yuta<sup>1\*</sup> ; SAITO, Akinori<sup>1</sup> ; SAKANOI, Takeshi<sup>2</sup> ; AKIYA, Yusuke<sup>1</sup> ; YAMAZAKI, Atsushi<sup>3</sup>

<sup>1</sup>Department of Geophysics, Graduate School of Science, Kyoto University, <sup>2</sup>Planetary Plasma and Atmospheric Research Center, Graduate School of Science, Tohoku University, <sup>3</sup>Institute of Space and Astronautical Science / Japan Aerospace Exploration Agency

Mesoscale wave-like structures in the mesospheric airglow were captured by an imaging observation from the International Space Station. Limb imaging observations with a digital single reflex camera provided us the horizontal structure of mesospheric airglow with unprecedentedly wide field-of-view(FOV). The FOV is 3,000 km width at the tangential point. Previous airglow observations captured small scale (10 - 400 km) structures by ground based airglow imagers and large scale (several 1,000s km) structures by satellite limb scanning. Our observation captured mesoscale wave-like structure, whose wavelength is about 1,000 km, and filled the gap of the previous observation. A data on August 26, 2015 shows wavelike oscillation in both peak intensity and layer height for OI, Na and OH airglow. Its wavelength is about 1,000 km and the wave existed region is about 2,500 x 3,000 km size over the Indian ocean and Australia. In the presentation, the cause of this structure will be discussed.

Keywords: airglow, gravity wave