

## 2011-2014年の福島県内の新田川における放射性セシウムの移行挙動 Transport of radiocesium in the Niida River, Fukuhsima Prefecture in 2011-2014

長尾 誠也<sup>1\*</sup>; 田堂 修<sup>1</sup>; 上村 輝宙<sup>1</sup>; 金森 正樹<sup>1</sup>; 桐島 陽<sup>2</sup>; 宮田 佳樹<sup>1</sup>; 落合 伸也<sup>1</sup>; 山本 政儀<sup>1</sup>  
NAGAO, Seiya<sup>1\*</sup>; TADO, Shu<sup>1</sup>; UEMURA, Teruaki<sup>1</sup>; KANAMORI, Masaki<sup>1</sup>; KIRISHIMA, Akira<sup>2</sup>; MIYATA, Yoshiki<sup>1</sup>; OCHIAI, Shinya<sup>1</sup>; YAMAMOTO, Masayoshi<sup>1</sup>

<sup>1</sup> 金沢大学, <sup>2</sup> 東北大学

<sup>1</sup>Kanazawa University, <sup>2</sup>Tohoku University

Surface deposition of <sup>134</sup>Cs and <sup>137</sup>Cs reveals considerable external radioactivity above 3000k Bq/m<sup>2</sup> in a zone extending northwest from the NPP after the Fukushima Daiichi Nuclear Power Plant (NPP) accident. Therefore, it is important to elucidate the short-term to long-term impacts of the accident on ecosystems of river watershed environments. This study investigated the transport of <sup>134</sup>Cs and <sup>137</sup>Cs in a small river, Niida River running through Iidate Viledge, in Fukushima Prefecture, Japan at normal and high flow conditions during 2011-2014.

Field experiments were conducted at a fixed station (Kinouchi bridge) in the lower Niida River during the period of May 2011-December 2014. The 20 L of surface river water samples were collected at the station using buckets. The radioactivity of <sup>134</sup>Cs and <sup>137</sup>Cs in the river waters before and after the filtration was measured with gamma-ray spectrometry using ammonium molybdophosphate (AMP)/Cs compound method.

Total radioactivity of <sup>137</sup>Cs (dissolved and particulate phases) in the river waters ranged from 0.11 to 4.18 Bq/L during May 2011-December 2014. Highest value was found in May 2011. Total <sup>137</sup>Cs radioactivity indicates the decreasing trend with increasing time at normal flow condition. However the higher radioactivity was observed after rain events. The <sup>137</sup>Cs radioactivity increased by 1.83 Bq/L after the heavy rain event by Typhoon Guchol in June 2012, and 1.68 Bq/L by Typhoon Jelawat in October 2012.

Percentage of <sup>137</sup>Cs associated with riverine suspended solids was 47-48% at normal flow condition in July and September 2011, but after December 2011 ranged from 75 to 93% at normal flow condition and 86-91% at high flow condition due to rain events. The radioactivity of particulate phase of <sup>137</sup>Cs ranged from 20 to 42 Bq/g-riverine suspended solids. Rain events are mainly contributed to the transport of radiocesium in the Niida River.

キーワード: Cs-134, Cs-137, 降雨イベント, 河川懸濁粒子

Keywords: Cs-134, Cs-137, rain event, riverine suspended solids