

## ガーナビリミアングリーンストーン帯ケープスリーポイント地域の地質構造及び層序の復元と堆積年代について

The stratigraphy and a new age constraint of the volcanoclastic sequence of Birimian in Cape Three Points, Ashanti belt, southwest Ghana

\*吉丸 慧<sup>1</sup>、清川 昌一<sup>1</sup>、伊藤 孝<sup>2</sup>、池原 実<sup>3</sup>、堀江 憲路<sup>4</sup>、竹原 真美<sup>4</sup>、佐野 貴司<sup>5</sup>、Nyame, K. Frank<sup>6</sup>、Tetteh, M George<sup>7</sup>

\*Satoshi Yoshimaru<sup>1</sup>, Shoichi Kiyokawa<sup>1</sup>, Takashi Ito<sup>2</sup>, Minoru Ikehara<sup>3</sup>, Kenji Horie<sup>4</sup>, Mami Takehara<sup>4</sup>, Takashi Sano<sup>5</sup>, Frank K Nyame<sup>6</sup>, George M Tetteh<sup>7</sup>

1.九州大学理学府地球惑星科学専攻、2.茨城大学教育学部、3.高知大学、4.国立極地研究所、5.国立科学博物館、6.ガーナ大学、7.タルクワ鉱山技術大学

1.Department of Earth and Planetary Science, Kyushu University, 2.Ibaraki University, 3.Kochi University, 4.National Institute of Polar Research, 5.National Museum of Nature and Science, 6.University of Ghana, 7.University of Mining and Technology

Paleoproterozoic Birimian greenstone belt, extending from eastern Guinea to western Niger, is located in southern part of West Africa Craton. The geological structure of the Ashanti belt occupying most of western Ghana, especially gold mines, has been revealed a little by scientific research. At the Cape Three Points area, there is no previous report about change in environment of deep sea floor during paleoproterozoic period. We focused on environment and bioactivities during deposition of protolith of the Birimian greenstones at this region. As a preliminary study, we report the stratigraphy, structures of volcanic and volcanoclastic sequence, the depositional settings and age at this region.

Ashanti belt, striking NE-SW and occupying most of western Ghana, is composed of mainly basalts or andesites, volcanoclastic rocks and belt type or non-belt type granitoids, and is uncomfortably overlain by acidic volcanoclastics and gold bearing conglomerates. The maximum depositional age of the cap of Birimian rocks is  $2154 \pm 2$  Ma (U-Pb zircon: Oberthür et al., 1998) and the youngest age of the intruded rock into Birimian volcanic in this region is  $2174 \pm 2$  Ma (U-Pb zircon: Oberthür et al., 1998).

In the Cape Three Points region facing the Gulf of Guinea, very thick volcanoclastic sequences are present in succession over 4000 m and about 1000 m-thick of stratigraphy of the study sites is reconstructed by detailed field investigations. The layers are running N-S, most of them dip 60-80 to the east. The stratigraphy shows fining upward. All of these sediments were affected by greenschist facies metamorphism, and thus minerals of amphiboles, plagioclase, chlorite and epidote are very common, while quartz grains are not so popular member here. The  $TiO_2/Al_2O_3$  ratio obtained from chromites with EPMA in basaltic rocks indicates that these rocks produced in the volcanic arc system. Trace elements compositions of whole-rock compositions tend to show low concentration of Nb, and high concentration of LREEs. These facts also indicate that they are derived from volcanic arc region. Those thick fining-upward volcanoclastic sequence, chemical compositions may be suggest that the Cape Three Points area was formed on the mid-deep sea floor beside an oceanic volcanic arc.

We obtained new age information from the porphyritic dyke with several meters in length and <1m in widths. This dyke is foliated and some minerals are sheared and tone off. Zircon grains collected from the dyke were measured by SHRIMP at NIPR, and yielded weighted mean  $^{204}Pb$ -corrected  $^{207}Pb/^{206}Pb$  age of  $2265.6 \pm 4.6$  Ma (95% confidence), which indicates that the volcanoclastic sequences deposited before  $2265.6 \pm 4.6$  Ma and deformed after that. Only four detrital zircons are dated around 2260 Ma in this site and these age groups are the oldest in the Ghanaian greenstone terrane (Loh and Hiedes

1992). Therefore, there is possibility that the early volcanic activity history of the Ghana Birimian remains at Cape Three Points.

キーワード：ガーナビリミアングリーンストーンベルト、古原生代、火山砕屑岩

Keywords: Ghana Birimian greenstone belt, paleoproterozoic, volcaniclastics