

Correlation-study about the ambient dose rate and the weather conditions

*Masato Furuya¹, Yuko Hatano¹, Tomoo Aoyama², Yasuhito Igarashi³, Kazuyuki Kita⁴, Masahide Ishizuka⁵

1.Department of Risk Engineering, Faculty of Systems and Information Engineering, University of Tsukuba, 2.Center for Research in Isotopes and Environmental Dynamics, University of Tsukuba, 3.Meteorological Research Institute, 4.College of Science, Ibaraki University, 5.Faculty of Engineering, Kagawa University

The long-term radiation risks are believed to be heavily affected by the resuspension process. We therefore focus on the surface-atmosphere exchange process of released radioactive materials in this study. Radioactive materials were deposited on the soil and float in the air, and such complicated process are influenced by the weather conditions deeply. We need to reveal the correlation between the weather conditions and the ambient dose rate.

In this study, we study the correlation between the weather conditions and the ambient dose rate with the correction of the decrease due to the radioactive decay. We found that there is a negative correlation between the ambient dose rate and the soil water content by the correlation coefficient. Using this result, we reconstruct the ambient dose rate from the weather conditions by the multiple regression analysis and found that the reconstructed data agree with the observation very well. Using Kalman filter, which can be sequentially updates the state estimate, we obtained such a good agreement.

Keywords: Fukushima Accident, Radioactive nuclides, Karman Filter