

## 太陽系小天体におけるホルムアルデヒドとアンモニアの化学

## Chemistry of formaldehyde and ammonia in the Solar System small bodies

\*癸生川 陽子<sup>1</sup>、三澤 柊介<sup>1</sup>、河合 純<sup>1</sup>、三田 肇<sup>2</sup>、南部 啓太<sup>3</sup>、大内 貴仁<sup>3</sup>、村松 康司<sup>3</sup>、橘 省吾<sup>4</sup>、小林 憲正<sup>1</sup>

\*Yoko Kebukawa<sup>1</sup>, Shusuke Misawa<sup>1</sup>, Jun Kawai<sup>1</sup>, Hajime Mita<sup>2</sup>, Keita Nanbu<sup>3</sup>, Takahito Ouchi<sup>3</sup>, Yasuji Muramatsu<sup>3</sup>, Shogo Tachibana<sup>4</sup>, Kensei Kobayashi<sup>1</sup>

1.横浜国立大学、2.福岡工業大学、3.兵庫県立大学、4.北海道大学

1.Yokohama National University, 2.Fukuoka Institute of Technology, 3.University of Hyogo, 4.Hokkaido University

Formaldehyde and ammonia is ubiquitous in the universe. Comets contain up to 4% H<sub>2</sub>CO and up to 1.5% NH<sub>3</sub> relative to H<sub>2</sub>O, and interstellar medium (ISM) contains up to 8% H<sub>2</sub>CO and up to 80% NH<sub>3</sub> relative to H<sub>2</sub>O [1]. Cody et al. [2] proposed the formation scenario of insoluble organic matter (IOM) in chondritic meteorites, in the presence of liquid water starting with formaldehyde and glycolaldehyde that is the simplest sugar produced by two H<sub>2</sub>CO molecules. We have conducted further experimental studies and showed that the presence of ammonia enhanced the IOM like organic solid formation via formose reaction followed by carbonization [3]. These studies were focusing on the solid materials, and now we study the liquid phase of the products with various analytical methods including electrospray ionization mass spectrometry (ESI-MS), X-ray absorption near edge structure (XANES), infrared spectroscopy and amino acid analyses using high performance liquid chromatography (HPLC).

Each starting solution contained 1mL water with 2 mmol formaldehyde, 1 mmol glycolaldehyde, 0.4 mmol ammonia (equivalent to H<sub>2</sub>O : C : N = 100 : 7.2 : 0.72) with catalytic amount of Ca(OH)<sub>2</sub>, and was sealed in a glass tube, then isothermally heated at 90 degrees C up to 250 degrees C. XANES and FTIR analyses showed that aromatic or olefinic C=C bond abundance increased with temperature in the soluble fractions. This indicates that the insoluble residues precipitate as a result of increase in the hydrophobic moieties in the products as the reaction proceeds. This is consistent with the previous results that the amount of insoluble fractions (organic solids) increases with temperature [3]. Acid hydrolysis of the solutions produced various amino acids up to four carbons. Alanine abundance was larger than glycine, and may indicate high abundance of methyl (-CH<sub>3</sub>) or methylene (-CH<sub>2</sub>-) substitutions in the amino acid precursor molecules. ESI-MS results suggested that various carbohydrates (CHO molecules) and these with nitrogen containing substitutions. This is somewhat consistent with ultrahigh-resolution ESI-MS analysis of the Murchison meteorite extract that shows various CHO and CHNO molecules [4], although only formaldehyde and ammonia chemistry cannot explain all of the diverse molecules found in this meteorite.

## References:

- [1] Charnley, S. B. and Rodgers, S. D. (2008) *Space Science Reviews*, 138, 59-73.
- [2] Cody, G. D., et al. (2011) *Proceedings of the National Academy of Sciences of the United States of America*, 108, 19171-19176.
- [3] Kebukawa, Y., et al. (2013) *The Astrophysical Journal*, 771, 19.
- [4] Schmitt-Kopplin, P., et al. (2010) *Proceedings of the National Academy of Sciences of the United States of America*, 107, 2763-2768.

キーワード：隕石、小惑星、有機物

Keywords: Meteorites, Asteroids, Organic matter