

## Broadband noise and associated electron heating observed by Kaguya around the Moon in the solar wind

\*Yasunori Tsugawa<sup>1</sup>, Yuto Katoh<sup>2</sup>, Naoki Terada<sup>2</sup>, Shinobu Machida<sup>1</sup>, Hideo Tsunakawa<sup>3</sup>, Futoshi Takahashi<sup>4</sup>, Hidetoshi Shibuya<sup>5</sup>, Hisayoshi Shimizu<sup>6</sup>, Masaki Matsushima<sup>3</sup>, Yoshifumi Saito<sup>7</sup>, Shoichiro Yokota<sup>7</sup>, Masaki N Nishino<sup>1</sup>

1.Institute for Space-Earth Environmental Research, Nagoya University, 2.Graduate School of Science, Tohoku University, 3.Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 4.Faculty of Sciences, Kyushu University, 5.Department of Earth and Environmental Sciences, Graduate School of Science and Technology, Kumamoto University, 6.Earthquake Research Institute, University of Tokyo, 7.Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency

Broadband electromagnetic noise in the frequency range up to  $\sim 10$  Hz has been detected around the Moon at  $\sim 100$  km altitude [Halekas et al., 2008; Nakagawa et al., 2011; Tsugawa et al., 2012]. Halekas et al. [2008] suggested that the waves are associated with electron energizations and are basically generated through the interaction between the solar wind plasma and crustal magnetic field. Nakagawa et al. [2011] studied the characteristics of the broadband waves by considering properties of whistler-mode waves propagating in the solar wind frame of reference. Tsugawa et al. [2012] showed that the statistical distributions of the intense noise are clearly located at the magnetic anomalies. While they discussed the possible generation process of the waves through resonant or non-resonant instability by ions reflected from the lunar surface, details of the generation process of the waves have not been clarified yet.

We analyzed the broadband noise observed by Kaguya statistically, and suggest that the absolute condition to observe the noise at altitudes  $\sim 100$  km are 1) the spacecraft is connected to the Moon through the magnetic field, and 2) the solar wind ions are reflected considerably in the connected region on the Moon. The fluxes of reflected ions depend on the solar wind parameters and the magnetisms of the lunar crusts. In a usual solar wind condition (roughly the dynamic pressure  $< 2$  nPa), the second condition is mostly satisfied above the magnetic anomalies. In the solar wind with larger density and faster speed than usual (roughly the dynamic pressure  $> 2$  nPa), the second condition can be satisfied above not only magnetic anomalies but also unmagnetized surface. Electrons are often energized perpendicular to the ambient magnetic field or isotropically in association with the noise and reflected ions. The electron heating above the lunar magnetic anomalies are also associated with the broadband electrostatic noise in the frequency range up to  $\sim 10$  kHz [Kasahara et al., 2011]. Their correlation is suggested in analogous to the transverse ion acceleration due to broadband extremely low frequency noise in the Earth's auroral region [e.g., Andre et al., 1998].