

Ultrahigh-pressure polymorphism in GeO₂ glass: implications for structure of magma at the core-mantle boundary

*Yoshio Kono¹, Curtis Kenney-Benson¹, Daijo Ikuta¹, Yuki Shibazaki², Yanbin Wang³, Guoyin Shen¹

1.Carnegie Institution of Washington, 2.Tohoku University, 3.The University of Chicago

Silicate magma at the core-mantle boundary is one of the most important components in understanding nature and evolution of the Earth's deep interior. However, structure and properties of silicate magmas at the pressure condition of the core-mantle boundary remain poorly understood, because of experimental challenges. Pioneering works by Murakami and Bass (2010; 2011) showed a kink in the pressure dependence of shear-wave velocity in SiO₂ and MgSiO₃ glasses around 130-140 GPa, which was interpreted as evidence of ultrahigh pressure structural transition. However, no structural information is available under such high pressures. Here we show new experimental evidence of ultrahigh pressure structural transition in GeO₂ glass with Ge-O coordination number (CN) significantly greater than 6, investigated using a newly developed double-stage large volume cell combined with multi-angle energy dispersive X-ray diffraction technique for in situ amorphous structure measurement. The Ge-O coordination number (CN) is found to remain constant at ~6 between 22.6 and 37.9 GPa. At higher pressures, CN begins to increase rapidly to 6.4 at 49.4 GPa and reaches 7.4 at 91.7 GPa. The structural change to CN higher than 6 is closely associated with the change in oxygen packing fraction (OPF). This transformation begins when the OPF in GeO₂ glass is close to the maximal dense packing state (the Kepler conjecture= ~ 0.74), which provides new insights into structural changes in network-forming glasses and liquids with CN higher than 6 at ultrahigh pressure conditions. For example, extrapolation of OPF-pressure trend in SiO₂ glass shows that OPF of SiO₂ glass reaches to 0.74 around 108 GPa, where structural change to CN higher than 6 is expected. The data imply that silicate magma at the core-mantle boundary may possess ultrahigh-pressure structure with CN higher than 6.

References

- Murakami, M., & Bass, J. D. (2010). Spectroscopic evidence for ultrahigh-pressure polymorphism in SiO₂ glass. *Physical review letters*, 104(2), 025504.
- Murakami, M., & Bass, J. D. (2011). Evidence of denser MgSiO₃ glass above 133 gigapascal (GPa) and implications for remnants of ultradense silicate melt from a deep magma ocean. *Proceedings of the National Academy of Sciences*, 108(42), 17286-17289.

Keywords: high pressure, magma, core-mantle boundary