

Accounting for the observation error correlation in data assimilation

*寺崎 康児¹、三好 建正¹

*Koji Terasaki¹, Takemasa Miyoshi¹

1. 理化学研究所計算科学研究機構

1. RIKEN Advanced Institute for Computational Science

It is known that some observations such as satellite radiances have the spatial and inter-channel error correlations. The current data assimilation systems in the operational centers mostly neglect the observation error correlations. It is important to account for the observation error correlations to effectively draw the information from the observation “BigData.”

In this study, we developed a method to account for the observation error correlations in the local ensemble transform Kalman filter (LETKF: Hunt et al. 2007), and performed idealized experiments with the Lorenz-96 model (Lorenz et al. 1998) and the non-hydrostatic icosahedral atmospheric model (NICAM: Satoh et al. 2013). The condition number of the observation error covariance matrix (**R** matrix), or the ratio of the maximum and minimum eigenvalues, is essential for the stable performance of the LETKF. Reconditioning can make the **R** matrix well-conditioned by adding a small constant to all the diagonal terms of the **R** matrix. We examined that the reconditioning not only stabilizes the LETKF but also greatly improves the analysis accuracy by including the observation error correlations.

キーワード：データ同化、観測誤差相関、条件数、Reconditioning

Keywords: Data assimilation, Observation error correlation, Condition number, Reconditioning