

ベトナム・ハノイにおける小型計測器によるPM_{2.5}の長期・多点測定 Using small compact PM_{2.5} sensors, long term and multipoint measurements in Hanoi, Vietnam

*岡本 渉¹、松見 豊¹、関口 和彦²、藤野 毅²、三小田 憲史²、熊谷貴 美代³、藤谷 雄二⁴、Thuy Ly Bich⁵、Dung Nghiem Trung⁵、Thuy Nguyen Thi Thu⁶、Thuy Pham Chau⁷、梶井 克純⁸、坂本 陽介⁸
*OKAMOTO WATARU¹, Yutaka Matsumi¹, Kazuhiko Sekiguchi², Takeshi Fujino², Kenshi Sankoda², Kimiyo Kumagai³, Yuji Fujitani⁴, Ly Bich Thuy⁵, Nghiem Trung Dung⁵, Nguyen Thi Thu Thuy⁶, Pham Chau Thuy⁷, Yoshizumi Kajii⁸, Yousuke Sakamoto⁸

1. 宇宙地球環境研究所 名古屋大学、2. 埼玉大学、3. 群馬県衛生環境研究所、4. 国立環境研究所、5. ハノイ工科大学、6. タイグエン工科大学、7. ベトナム国立農業大学、8. 京都大学

1. ISEE Nagoya University, 2. Saitama University, 3. Gunma Prefecture Sanitary Environment Laboratory, 4. National Institute for Environmental Studies, 5. Hanoi University of Science and Technology, 6. Thai Nguyen University of Technology, 7. Vietnam National University of Agriculture, 8. Kyoto University

A year round observation of PM_{2.5} mass concentrations was conducted in Hanoi University of Science and Technology (HUST) from July 2016 to now using newly developed highly sensitive sensors. And also from October 2017 to now, we have installed the PM_{2.5} sensors in Thai Nguyen University of Technology (TNUT) and Vietnam National University of Agriculture (VNUA). Hourly PM_{2.5} concentration data in HUST agreed well with corresponding data obtained with a beta attenuation monitor, located 3 km away. The monthly variation showed that PM_{2.5} levels were high in the dry season, particularly in December (median = 62 mg m⁻³) and low in the rainy season, particularly in June and July (medians = 19 mg m⁻³). PM_{2.5} haze episodes with levels higher than 100 mg m⁻³ were observed 13 times during the dry season (October 2016–March 2017). These episodes may link to meteorological conditions, because the daily PM_{2.5} and CO levels increased several days after a cold surge for most of the PM_{2.5} haze episodes. Two episode events were investigated further in higher temporal resolution. For both events, PM_{2.5} peaks appeared at midnight for some days after a cold surge, while CO levels often increased at rush hours. For first event, short peaks of PM_{2.5} (duration of about 1 h) with fast rates of increase of ~5 mg m⁻³ min⁻¹ were observed. This highlights the importance of high temporal resolution of PM_{2.5} sensors in assessing of health effect. For second event, a broad peak (duration of one and half days) of PM_{2.5} with gradual increase was observed together with high CO levels. This research reveals for first time characteristics of PM_{2.5} haze episode in Hanoi that should be studied further to develop an effective air pollution management in this city. This study is supported by JSPS Kakenhi (17H04483).

キーワード：PM_{2.5}測定、小型測定器、空気の良さ、大都市、東南アジア

Keywords: PM_{2.5} measurement, compact sensor, air quality, Mega city, Southeast Asia

