

## Hindcasts of the 2016 Disruption of the Stratospheric Quasi-biennial Oscillation

\*渡辺 真吾<sup>1</sup>、Kevin Hamilton<sup>2</sup>、Scott Osprey<sup>3</sup>、河谷 芳雄<sup>1</sup>、西本 絵梨子<sup>1</sup>

\*Shingo Watanabe<sup>1</sup>, Kevin Hamilton<sup>2</sup>, Scott Osprey<sup>3</sup>, Yoshio Kawatani<sup>1</sup>, Eriko Nishimoto<sup>1</sup>

1. 国立研究開発法人海洋研究開発機構、2. International Pacific Research Center, Univ. of Hawaii、3. NCAS-Climate, Univ. of Oxford

1. Japan Agency for Marine-Earth Science and Technology, 2. International Pacific Research Center, Univ. of Hawaii, 3. NCAS-Climate, Univ. of Oxford

In early 2016 the quasi-biennial oscillation in tropical stratospheric winds was disrupted by an anomalous easterly jet centered at ~40 hPa, a development that was completely missed by all operational extended range weather forecast systems. This event and its predictability are investigated through ensemble hindcasts using a global model notable for its sophisticated representation of the upper atmosphere. Key to prediction of this event is simulating the slowly evolving mean winds in the winter subtropics that provide a waveguide for Rossby waves propagating from the winter hemisphere. Its association to the strong El Nino condition in 2015 is also investigated by changing sea surface temperatures.

キーワード : disruption of quasi-biennial oscillation、hindcast、high-vertical resolution global climate model

Keywords: disruption of quasi-biennial oscillation, hindcast, high-vertical resolution global climate model