## Direct observation of elementary steps and quasi-liquid layers on ice surfaces by advanced optical microscopy

\*Ken Nagashima<sup>1</sup>, Gen Sazaki<sup>1</sup>, Ken-ichiro Murata<sup>1</sup>, Yoshinori Furukawa<sup>1</sup>, Tetsuya Hama<sup>1</sup>

1. The Institute of Low Temperature Science, Hokkaido University

Ice is one of the most abundant crystals on the earth, and hence the molecular-level understanding of ice crystal surfaces holds the key to unlocking the secrets of a number of fields. We and Olympus Engineering Co., Ltd. have developed laser confocal microscopy combined with differential interference contrast microscopy (LCM-DIM), by which we succeeded in the direct visualization of 0.37-nm-thick elementary steps on ice for the first time [1]. Subsequently, the direct observations on ice basal faces revealed the various features of elementary steps [2-4].

On the other hand, we could also visualize the quasi-liquid layers (QLLs) on ice crystal surfaces [5], which are covered with thin liquid layers even below the melting point (0°C). The direct observations of QLLs revealed the appearance of two types of QLLs with different morphologies [5,6], the appearance temperatures and partial pressure of water vapor [6-8], the inducement of the formation of QLLs by strain [9], and the characteristic velocities of QLLs [10].

In addition, we also found that atmospheric acidic gas (hydrogen chloride gas) strongly induced the appearances of droplets on ice surfaces (further details will be presented in "A-ASO6: Atmospheric Chemistry" session). The droplets were observed in the temperature range of -15.0  $^{\sim}$  -1.5 °C, where no QLL appears in the absence of HCl gas [11]. The HCl induced droplets were embedded into ice crystals by growth of ice crystals [12]. These results show the possibility that ice crystals can store large amount of gas components as fluid inclusions.

- [1] Sazaki et al. (2010) PNAS 107, 19702.
- [2] Sazaki et al. (2014) Cryst. Growth Des. 14, 2133.
- [3] Asakawa et al. (2014) Cryst. Growth Des. 14, 3210.
- [4] Inomata et al. (2018) Cryst. Growth Des. 18, 786.
- [5] Sazaki et al. (2012) PNAS 109, 1052.
- [6] Murata et al. (2016) PNAS 113, E6741.
- [7] Asakawa et al. (2015) Cryst. Growth Des. 15, 3339.
- [8] Asakawa et al. (2015) PNAS 113, 1749.
- [9] Sazaki et al. (2013) Cryst. Growth Des. 13, 1761.
- [10] Murata et al. (2015) Phys. Rev. Lett. 115, 256103.
- [11] Nagashima et al. (2016) Cryst. Growth Des. 16, 2225.
- [12] Nagashima et al., submitted.

Keywords: Ice, Surface melting, Quasi-liquid layer, Molecular step, Advanced optical microscopy