

Structure from Motion技術を用いたネパールヒマラヤ・トランバウ氷河 における2007-2017年の表面標高変化

Surface elevation change from 2007 to 2017 using Structure from Motion at the Trambau glacier, Nepal

*佐藤 洋太¹、藤田 耕史¹、井上 公²、對馬 あかね³、砂子 宗次朗¹、坂井 亜規子¹

*Yota Sato¹, Koji Fujita¹, Hiroshi Inoue², Akane Tsushima³, Sojiro Sunako¹, Akiko Sakai¹

1. 名古屋大学環境学研究科、2. 防災科学研究所、3. 総合地球環境学研究所

1. Graduate School of Environmental Studies, Nagoya University, 2. National Research Institute for Earth Science and Disaster Resilience, 3. Research Institute for Humanity and Nature

Alpine glaciers are good indicator of climate change, and recent shrinkage of glaciers contribute to the sea level rise. In addition to that, shrinkage of glaciers is increasing potential exposure to glacier lake outburst floods (GLOFs) in the Himalaya (e.g. Benn et al., 2012). Therefore, monitoring mass balance of Himalayan glaciers is important to assess glacier fluctuation.

Digital elevation model (DEM) based on satellite imageries have been analyzed to estimate fluctuation of glacier mass balance. This method permit getting data covering wide area but it's difficult to obtain high resolution surface elevation changes. Some previous studies suggested that ice cliffs and ponds have significant roles in mass loss of debris-covered glaciers (e.g. Buri et al., 2016). Almost DEMs and orthomosaic based on satellite images are too coarse to extract them. However, Structure from Motion (SfM) technology based on aerial photographs has been developed in the last few years. This method can create 3D model from photographs and it enables generating DEMs and orthomosaic image with high resolution. It allows us to identify the small structures on glaciers. Aerial photograph of glacier surface can be collected relatively easy and they are not expensive comparing to other remote sensing data. Thus, some studies combined aerial photogrammetry and SfM technology have been carried out in the Nepal Himalayan region (e.g. Immerzeel et al., 2014).

In this study, we created high resolution DEMs from 10 years interval aerial photographs by using SfM technology at Trambau glacier in the Nepal Himalaya. These photographs were taken by traditional airborne (plane) in 2007 and taken by handmade fixed-wing UAV in 2017. We also conducted GPS measurement in 2017 post-monsoon season to obtain elevation and coordinate data for georeference and to evaluate the DEMs accuracy. Consequently, we calculated spatial elevation changes from the two DEMs in the Trambau glacier.

We are going to talk about results of the DEMs accuracy and spatial surface elevation changes at this area.

References

Benn et al., 2012: Earth Sci. Rev. 114 (1-2), 156-174.

Buri et al., 2016: Ann. Glaciol. 57 (71), 199-211.

Immerzeel et al., 2014: Remote Sens. Environ. 150, 93-103.

キーワード：氷河、ヒマラヤ、SfM、UAV、リモートセンシング、空撮

Keywords: Glacier, Himalaya, Structure from Motion, Unmanned Aerial Vehicle (UAV), Remote Sensing, Aerial photogrammetry