

走査型電子顕微鏡を使ったグリーンランド氷床北西部SIGMA-Dアイスコア中の鉱物組成変動の解明

Variations in composition of mineral dust in an ice core obtained from Northwestern Greenland Ice Sheet

*永塚 尚子¹、東 久美子^{1,2}、本山 秀明^{1,2}、的場 澄人³、藤田 耕史⁴、山崎 哲秀⁵、大沼 友貴彦⁶、箕輪 昌紘³、青木 輝夫⁷、平林 幹啓¹、Dallmayr Remi⁸

*Naoko Nagatsuka¹, Kumiko Goto-Azuma^{1,2}, Hideaki Motoyama^{1,2}, Sumito Matoba³, Koji Fujita⁴, Tetsuhide Yamasaki⁵, Yukihiko Onuma⁶, Masahiro Minowa³, Teruo Aoki⁷, Motohiro Hirabayashi¹, Dallmayr Remi⁸

1. 国立極地研究所、2. 総合研究大学院大学、3. 北海道大学低温科学研究所、4. 名古屋大学大学院環境学研究科、5. アバソナット、6. 東京大学生産技術研究所、7. 岡山大学自然科学研究科、8. AWI

1. National Institute of Polar Research, 2. SOKENDAI (The Graduate University for Advanced Studies), 3. Institute of Low Temperature Science, Hokkaido University, 4. Graduate School of Environmental Studies, Nagoya University, 5. Avangnaq, 6. Institute of Industrial Science, University of Tokyo, 7. Graduate School of Natural Science and Technology, Okayama University, 8. AWI

Snow and ice on glaciers and the ice sheet in the Arctic contain windblown mineral dust derived from local sediments as well as distant deserts. Dust deposited on the ice sheet in the past can be obtained by ice core drilling, and the variations in its sources and transportation processes can be reconstructed by particle analysis of ice cores. In this study, we analyzed morphology and surface chemistry of mineral dust particles in an ice core drilled in Northwest Greenland with Scanning Electron Microscope (SEM, QUANTA FEG 450) and Energy Dispersive X-ray Spectrometer (EDS).

The ice core was drilled at the SIGMA-D site (N77°64', W59°12') of 2100 m a.s.l. in 2014. The length is 222.72 m and the estimated age at 113 m depth is 350 years before present. The ice samples were collected every five years in plastic bottles and freeze dried on a polycarbonate filter to concentrate micro-particles. Then, the filter was coated with platinum (Pt) for SEM analysis.

The SEM observation revealed that the mean size of mineral dust in the SIGMA-D ice core ranged from 1-3 μm , which is similar to that of the other Greenland ice core dust that seems to be derived from distant deserts. This suggests that the SIGMA-D ice core contained mainly long-range transported wind-blown mineral dust. The EDS analysis also revealed the ice core contained mainly silicate minerals, especially clay minerals, that showed compositional variation among the samples. For example, Quartz and Feldspar contents were three times higher in 1935-1950 compared with the other periods. Furthermore, the variation trend also differed among the minerals. The trend of micas/illite, chlorite, and feldspar were similar to each other, but different from kaolinite. The two type of minerals were formed by different weathering processes. These results indicate that the ice core dust was likely derived from different sources in different period.

キーワード：アイスコア、グリーンランド、走査型電子顕微鏡、鉱物組成

Keywords: ice core, Greenland, Scanning Electron Microscope (SEM), mineral composition