Plans for developing an Earth system model with CH₄ dynamics

*井上 忠雄¹、羽島 知洋¹、伊藤 昭彦²
*Tadao Inoue¹, Tomohiro Hajima¹, Akihiko Ito²

- 1. 国立研究開発法人海洋研究開発機構、2. 国立研究開発法人国立環境研究所
- 1. Japan Agency for Marine-Earth Science and Technology, 2. National Institute for Environmental Studies

Methane ($\mathrm{CH_4}$) is one of the important GHGs for climate prediction due to its relatively high radiative forcing next to that of $\mathrm{CO_2}$, and now it is necessary to pay more attention to global $\mathrm{CH_4}$ dynamics after Paris agreement. Under the control of $\mathrm{CO_2}$ emission to keep a global temperature rise this century below 2 degrees Celsius above pre-industrial level, concentration of $\mathrm{CH_4}$ in the atmosphere will be more influential to climate change. Furthermore, the atmospheric life time of $\mathrm{CH_4}$ is around 9 years, it could be a good target for climate change mitigation. However, current Earth system models (ESMs) cannot project $\mathrm{CH_4}$ concentration with fully coupled manner: some models can predict atmospheric $\mathrm{CH_4}$ concentration but its predicted concentration is not based on the anthropogenic/natural $\mathrm{CH_4}$ emission under climate interactions. In order to make climate projection with fully coupled $\mathrm{CH_4}$ processes, we have started to couple $\mathrm{CH_4}$ dynamics into our ESM, by introducing atmospheric chemistry model and the terrestrial $\mathrm{CH_4}$ emission scheme into an ESM. In this presentation, brief introduction and the progress of our ESM development will be made, and plans for application of the ESM to $\mathrm{CH_4}$ -related simulations will be shown.

キーワード:気候変動、メタン、地球システムモデル

Keywords: Climate Change, Methane(CH4), Eearth System Model(ESM)