

Variation in coral calcifying fluid pH across coral habitable latitude

*田中 健太郎¹、Holcomb Michael^{2,3}、渡邊 剛⁴、McCulloch Malcolm^{2,3}

*Kentaro Tanaka¹, Michael Holcomb^{2,3}, Tsuyoshi Watanabe⁴, Malcolm McCulloch^{2,3}

1. 東京大学 大気海洋研究所、2. School of Earth and Environment, The University of Western Australia、3. ARC Centre of Excellence for Coral Reef Studies, The University of Western Australia、4. 北海道大学大学院理学院自然史科学専攻

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. School of Earth and Environment, The University of Western Australia, 3. ARC Centre of Excellence for Coral Reef Studies, The University of Western Australia, 4. Department of Natural History Sciences, Hokkaido University

Scleractinian corals are distributed throughout the coastal area across the tropical to the temperate oceans. Their aragonite skeletons contribute to reef development and the reef ecosystem. The rate of coral calcification is influenced by several factors (temperature, pH, and light intensity), which differs among coral habitats, generally being higher in the tropical zone than in the temperate zone. In addition, current global climate changes (global warming and ocean acidification) possibly impact coral calcification, and thus coral reef formation and ecosystem. It is essential to understand the main factors controlling coral calcification in order to predict changes in coral calcification in the future associated with changing coastal environments.

Since the accretion of coral skeletons occurs within the calcifying fluid (CF) between coral soft tissues and hard skeletons, the chemical composition of the CF, such as pH and aragonite saturation state (Ω_{ara}), is likely to influence coral calcification. In this study, the variation in pH and Ω_{ara} in CF of the reef-building corals collected from a wide latitudinal range (from $\sim 7^{\circ}\text{N}$ to 34°N) are derived using the geochemical tools to understand the contribution of environment to the chemical composition in the CF.

The pH and Ω_{ara} in the CF were derived from boron isotopes and boron concentration in the coral skeletons. Specimens were collected from the tropical ($\sim 7^{\circ}\text{N}$) to the temperate ocean ($\sim 34^{\circ}\text{N}$) and hence experienced different temperatures (annual average between $\sim 22 - 29^{\circ}\text{C}$). The pH in the CF showed an increasing trend from lower to higher latitude with decreasing temperature. We found that Ω_{ara} in the CF ranges between 10-20 without a significant relation with latitude. This fact suggests corals can maintain Ω_{ara} in CF within similar range regardless of the difference in seawater environments among their habitats. Although Ω_{ara} is similar among corals, calcification rate is generally higher in tropical zone, implying that temperature mainly controls the coral calcification rate.

キーワード：サンゴ、バイオミネラリゼーション、pH、アラゴナイト飽和度

Keywords: coral, biomineralisation, pH, aragonite saturation state