

## Possibility of Nitrogen Emissions Evaluation from the Viewpoint of State-Scale Agricultural Product Production and Transport in Brazil

\*宮岡 邦任<sup>1</sup>、福岡 正人<sup>2</sup>、田瀬 則雄<sup>3</sup>、小野寺 真一<sup>2</sup>、齋藤 光代<sup>4</sup>、清水 裕太<sup>5</sup>、シロタ リカルド<sup>6</sup>  
\*Kunihide Miyaoka<sup>1</sup>, Masato Fukuoka<sup>2</sup>, Norio Tase<sup>3</sup>, Shin-ichi Onodera<sup>2</sup>, Mitsuyo Saito<sup>4</sup>, Yuta Shimizu<sup>5</sup>, Ricardo Shirota<sup>6</sup>

1. 三重大学教育学部、2. 広島大学大学院総合科学研究科、3. 茨城大学教育学部、4. 岡山大学大学院環境生命科学研究科、5. 農業・食品産業技術総合研究機構 西日本農業研究センター、6. サンパウロ大学農学部

1. Faculty of Education, Mie University, 2. Graduate School of Integrated Arts and Science, Hiroshima University, 3. Faculty of Education, Ibaraki University, 4. Graduate school of environmental and life science, Okayama University, 5. National Agriculture and Food Research Organization, Western Region Agricultural Research Center, 6. College of Agriculture, Sao Paulo University

In Brazil, the major exporting country of agricultural products, the production of sugar cane and soybean are increasing in recent years, because of growing global demand for bioethanol and for worldwide food situation. Regional disparities are remarkable among states, prices and labor costs are different as well. It is thought that these regional differences reflects in the cost per unit of sugar cane production. There is a possibility that the transportation cost can be offset when considering the total balance of personnel expenses, mechanization, energy costs (water supply and electricity), etc. even in areas far from consumers and ports. Also, depending on natural environment and cultivar varieties, there may be regional differences in fertilizer application

In this study, we estimate the amount of nitrogen emission per unit production amount, considering the different social and natural environments of nitrate contamination of groundwater by local fertilization. Based on the results, consider appropriate sugar cane cultivation form and nitrogen emission control for sustainable development. In this presentation, as the first stage of research, we discuss that may estimate nitrogen emissions in each state by considering socio-economic environment effect, changes in sugar cane production volume, distribution form, and distribution cost.

The population pyramid shows that many states in the south are types of developed countries, whereas in the north there are many states of the type of developing countries. The distribution of economic power by city shows that the economic power is high in the large cities of the south and its surrounding cities, whereas in the northern part the major cities such as Recife have some economic power, but there are relatively low in the many city. Sugar cane production areas are concentrated in Pernambuco State, Aragoas State, Minas Gerais State, Sao Paulo State, Mato Grosso do Sul State. The economic power of these states is high throughout the state in Sao Paulo State and Minas Gerais State in the southern part, but in the states in the northeastern region there is a big disparity within the state. Produced sugar cane is accumulated in Recife in the northeastern part and accumulated in Santos in the south. The cost of being transported from various parts of the southern part to Santos is not cheaper as the distance is shorter, and there is no proportional relation between distance and transportation cost.

Further, it is needs to collect the data, such as transportation costs and labor costs for each city.

Acknowledgement: This study has supported by JSPS KAKENHI Grant Number 16KT0033.

キーワード：ブラジル、サトウキビ、地域格差、輸送コスト、窒素排出量

Keywords: Brazil, sugar cane, regional disparities, transportation costs, nitrogen Emissions

