

The Formation Process of 3D normal-mode Energy Spectra using Numerical Simulation of Rossby Wave Breaking

*松信 匠¹、田中 博²

*Takumi Matsunobu¹, Hiroshi L Tanaka²

1. 筑波大学生命環境学群地球学類、2. 筑波大学計算科学研究センター

1. College of Geoscience, School of Life and Environmental Sciences, University of Tsukuba, 2. Center for Computational Sciences, University of Tsukuba

Energy spectra represented by a three-dimensional normal mode expansion follow the slope of $E = (p_s / g) c^2$. Rossby wave breaking plays an important role in making energy spectra. Tanaka and Watarai (1999) said wave breaking is caused by increasing the energy of noise waves. $E = (p_s / g) c^2$ is derived from the Garcia (1991)'s criterion $(dq/dy) < 0$, representing the local meridional gradient of potential vorticity becomes negative, in Tanaka et al. (2004).

In this study, we verified the criteria of saturation and breaking through time. First, we watched how the energy is transferred by each factor. The wave-wave interactions begin with the saturation of transportation from wavemaker to its harmonics. Second, the energetic relation among waves is researched. The local meridional gradient of potential vorticity becomes negative in PV contours as the energy becomes $E = (p_s / g) c^2$, but wave breaking isn't necessary occurred. The transportation to noise waves is excited by the saturation of wavemaker and harmonics. Wave breaking occurs when the energy of noise wave becomes comparable with the total energy of harmonics. The results agree with the Tanaka and Watarai (1999)'s criterion of wave breaking and the Tanaka et al. (2004)'s saturation criterion. Third, the baroclinic instability was injected under the fixed ratio. Transferring energy to the noise waves sets up the Rossby wave breaking and the structure of baroclinic instability defines the transportation of wave-wave interaction.

キーワード：ノーマルモードエネルギー論、3次元ノーマルモード、ロスビー砕波

Keywords: Normal-mode energetics, Energy spectrum, Rossby wave breaking