

## An Expanded Gas-Grain Model for Interstellar Glycine

\*鈴木 大輝<sup>1</sup>、Majumdar Liton<sup>2</sup>、大石 雅寿<sup>3</sup>、齋藤 正雄<sup>3</sup>、廣田 朋也<sup>3</sup>、Valentine Wakelam<sup>4</sup>

\*Taiki Suzuki<sup>1</sup>, Liton Majumdar<sup>2</sup>, Masatoshi Ohishi<sup>3</sup>, Masao Saito<sup>3</sup>, Tomoya Hirota<sup>3</sup>, Valentine Wakelam<sup>4</sup>

1. 自然科学研究機構アストロバイオロジーセンター、2. ジェット推進研究所、3. 国立天文台、4. ボルドー大学

1. National Institutes of Natural Sciences Astrobiology Center, 2. Jet Propulsion Laboratory, NASA, 3. National Astronomical Observatory of Japan, 4. Bordeaux University

The study of the chemical evolution of glycine in the interstellar medium is one of challenging topics in astrochemistry. Here, we present the chemical modeling of glycine in hot cores using the state-of-the-art three-phase chemical model NAUTILUS, which is focused on the latest glycine chemistry. For the formation process of glycine on the grain surface, we obtained consistent results with previous studies that glycine would be formed via the reactions of COOH with CH<sub>2</sub>NH<sub>2</sub>. However, we will report three important findings regarding the chemical evolution of interstellar glycine. First, with the experimentally obtained binding energy from the temperature programmed thermal desorption (TPD) experiment, large part of glycine was destroyed through the grain surface reactions with NH or CH<sub>3</sub>O radicals before it fully evaporates. As a result, the formation process in the gas phase is more important than thermal evaporation from grains. If this is the case, NH<sub>2</sub>OH and CH<sub>3</sub>COOH rather than CH<sub>3</sub>NH<sub>2</sub> and CH<sub>2</sub>NH would be the essential precursors to the gas phase glycine. Secondly, since the gas phase glycine will be quickly destroyed by positive ions or radicals, early evolutionary phase of the hot cores would be the preferable target for the future glycine surveys. Thirdly, we suggest the possibility that the suprathreshold hydrogen atoms can strongly accelerate the formation of COOH radicals from CO<sub>2</sub>, resulting in the dramatical increase of formation rate of glycine on grains. The efficiency of this process should be investigated in detail by theoretical and experimental studies in the future.

キーワード：星間化学、化学進化、グリシン

Keywords: Interstellar Chemistry, Chemical Evolution, Glycine