

## Characteristics of O and Si isotope ratios of zircons of silicic rocks from the Izu-Bonin-Mariana (IBM) arc and the Izu collision zone (ICZ)

\*Takayuki Ushikubo<sup>1</sup>, Kenichiro Tani<sup>2</sup>, R. Bastian Georg<sup>3</sup>

1. Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology, 2. Department of Geology and Paleontology, National Museum of Nature and Science, 3. Trent University Water Quality Center

The Izu-Bonin-Mariana (IBM) Arc is an active intra-oceanic arc for ~52 million years, and silicic crusts have been generated by subduction zone magmatism [e.g., 1,2]. The north end of the IBM arc has been colliding with the Honshu Arc at the Izu Collision Zone (ICZ) for ~15 million years and voluminous syn-collision granitic plutons are exposed in ICZ [e.g., 3]. Silicic crust formations by intra-oceanic crustal subduction and the following arc-arc collision are probably significant processes of continental crust formation during the early Earth history.

Elevated  $\delta^{18}\text{O}$  values (6.0–7.5‰, VSMOW) in Hadean and Archean zircons are interpreted as evidence for incorporation of low temperature aqueous alteration products to the zircon-forming magmas as early as 4.3 Ga [4,5]. Similarly, low  $\delta^{30}\text{Si}$  magmas can be caused by incorporation of severely weathered materials [6,7]. Oxygen and Si isotope ratios of zircons can be useful to investigate incorporation of low-temperature weathered material to their parent magmas. We established zircon standards, a kimberlite xenocryst KC-KLV-Zrc1 from Kaalvallei, South Africa ( $\delta^{18}\text{O}_{\text{VSMOW}}=5.43\pm 0.14\%$ ,  $\delta^{30}\text{Si}_{\text{NBS-28}}=-0.38\pm 0.19\%$ , 2SD) and a detrital zircon megacryst KC-MGK-Zrc2 from Mogok, Myanmar ( $\delta^{18}\text{O}_{\text{VSMOW}}=19.14\pm 0.14\%$ ,  $\delta^{30}\text{Si}_{\text{NBS-28}}=-0.18\pm 0.09\%$ , 2SD), for in situ O and Si isotope measurements by SIMS. And we performed O and Si isotope analyses of zircons from IBM Arc (6 granitic rocks and 1 gabbro, ~52 to 0.15 Ma) and ICZ (6 granitic rocks, 15.7 to 4.3 Ma) by SIMS. Typical spot-to-spot reproducibility (2SD) of SIMS measurements was  $\pm 0.25\%$  for  $\delta^{18}\text{O}$  and  $\pm 0.3\%$  for  $\delta^{30}\text{Si}$ , respectively. Part of isotope data have already reported in JpGU 2016 [8].

The zircon  $\delta^{18}\text{O}$  values of most IBM granitic rocks (4.9 to 5.2‰) are within the range of the mantle-like zircon value ( $5.3\pm 0.6\%$ , [4,9]) with one exception of the Omachi Seamount sample (7.0‰). Zircons from IBM fore-arc gabbro have slightly low values (4.6‰). In contrast, zircons from ICZ tend to have elevated  $\delta^{18}\text{O}$  values (up to 6.4‰), indicating incorporation of accreted sediments in the Honshu Arc. The  $\delta^{30}\text{Si}$  values of all measured zircons ( $-0.4\pm 0.3\%$ ) are consistent with the kimberlite zircon within analytical uncertainty ( $\sim \pm 0.3\%$ ). The present results suggest that incorporation of sediments to produce elevated  $\delta^{18}\text{O}$  magmas effectively occurs by the arc-arc collision process but production of low  $\delta^{30}\text{Si}$  magma require further accumulation and incorporation of matured sediments.

### References:

- [1] Ishizuka et al. (2011) *EPSL*, 306, 229-240.
- [2] Tani et al. (2015) *EPSL*, 424, 84-94.
- [3] Tani et al. (2010) *Geology*, 38, 215-218.
- [4] Valley et al. (2005) *Contrib. Mineral. Petrol.*, 150, 561-580.
- [5] Valley et al. (2015) *Am. Mineral.*, 100, 1355-1377.
- [6] Savage et al. (2014) *Lithos*, 190-191, 500-519.
- [7] Savage et al. (2016) Goldschmidt Conference abstract #2749.
- [8] Tani et al. (2016) JpGU abstract #SGL38-08.
- [9] Grimes et al. (2011) *Contrib. Mineral. Petrol.*, 161, 13-33.

Keywords: zircon, IBM arc, SIMS, oxygen isotope, silicon isotope