

# コマチアイトの硫黄同位体異常を用いた太古代マントル物質循環の研究

## The study on material cycle of Archean mantle: Sulfur isotope anomaly of komatiite

\*久保田 勇祐<sup>1</sup>、上野 雄一郎<sup>1,2,3</sup>、清水 健二<sup>2</sup>

\*Yusuke Kubota<sup>1</sup>, Yuichiro Ueno<sup>1,2,3</sup>, Kenji Shimizu<sup>2</sup>

1. 東京工業大学理学院地球惑星科学系、2. 海洋研究開発機構、3. 東京工業大学地球生命研究所

1. Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 2. Japan Agency for Marine-Earth Science and Technology, 3. Earth-Life Science Institute, Tokyo Institute of Technology

Multiple sulfur isotopes of a 2.7 Ga komatiite and related volcanics were measured to study recycled crustal material in Archean mantle. The measured komatiites are from Belingwe Greenstone Belt, Zimbabwe, which represent one of the best-preserved Archean komatiite in the world. The Belingwe volcanics has been classified into four groups: komatiite, komatiitic basalt, D-basalt and E-basalt. The results show that some komatiites and D-basalts exhibit clear S-MIF ( $-0.204\% < \Delta^{33}\text{S} < -0.115\%$ ), whereas all the other samples do not show S-MIF ( $-0.025\% < \Delta^{33}\text{S} < +0.035\%$ ). Based on petrological observation of the S-MIF-bearing samples, the sulfide minerals were crystallized at the time of eruption, but not introduced after the emplacement. The S-MIF signature only occurred in depleted rocks (i.e., komatiite and D-basalt), but not in enriched rocks (i.e., komatiitic basalt and E-basalt), suggesting that the S-MIF source were not assimilated from surrounding Archean crust at the time of eruption, but derived from the source mantle. The S-MIF bearing rocks also show lower initial  $^{238}\text{U}/^{204}\text{Pb}$ , higher  $\epsilon\text{Nd}$  and lower  $\epsilon\text{Sr}$  values than non-S-MIF bearing rocks. All these characteristics support that the source mantle contained the MIF sulfur materials. The S-MIF signature may have been partly erased during the crustal contamination process or represent heterogeneous source in the Archean mantle. In either case, the results of this study indicate that Archean mantle contained S-MIF bearing materials, which probably represent subducted crustal material. Irrespective to tectonic style, transfer of surface material into the mantle has already been operated at least older than 2.7 Ga.

キーワード：硫黄同位体、太古代、マントル物質循環

Keywords: Sulfur isotopes, Archean, Mantle material cycle