

Carbon-cycle perturbations in the deep lapetus during the Great Ordovician Biodiversification Event

*藤崎 渉¹、松井 洋平¹

*Wataru Fujisaki¹, Yohei Matsui¹

1. 国立研究開発法人海洋研究開発機構

1. Japan Agency for Marine-Earth Science and Technology

The Ordovician represents a period of major biospheric change in the Earth history, including a long-term biodiversification followed by a short-term extinction; the latter has been linked to a global cooling episode. The three-fold increase in biodiversity of throughout the Ordovician is known as the Great Ordovician Biodiversification Event (GOBE), which has been a focus of numerous paleo-environmental investigations in the fossiliferous shallow-marine strata deposited along continental margins. However, strata of deep-sea facies have not been focused in previous paleo-environmental studies, although they occupied extensive oceanic domains.

Deep-sea bedded cherts in the Ballantrae accretionary complex in SW Scotland is appropriate for checking the Ordovician extinction-related paleo-environmental information in the deep lapetus Ocean. To examine the perturbations in C-cycle in the deep oceans during the GOBE, we preliminary report the organic carbon isotopic records from the Middle Ordovician deep-sea cherts at Bennane Head, Ballantrae Complex, SW Scotland.

キーワード：オルドビス紀生物多様性イベント、有機炭素同位体比、深海チャート、スコットランド

Keywords: Great Ordovician Biodiversification Event (GOBE), organic carbon isotopes, deep-sea bedded cherts, Scotland