[EJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-AS Atmospheric Sciences, Meteorology & Atmospheric Environment

## [A-AS06]Atmospheric Chemistry

convener:Yoko Iwamoto(Graduate School of Biosphere Science, Hiroshima University), Tomoki Nakayama(Graduate School of Fisheries and Environmental Sciences, Nagasaki University), Sakae Toyoda(東京工業大学物質理工学院, 共同), Nawo Eguchi(Kyushu University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) This session provides a forum for the presentation of the broad spectrum of tropospheric and stratospheric chemistry, including various research topics (e.g., dynamical processes, air quality and climate), approaches (modeling, field measurements, remote sensing, and laboratory studies), and species (gas and aerosol). This session also provides an opportunity for discussing possible future collaboration with other research fields relevant to atmospheric chemistry.

## [AAS06-P03]Working standard gas saving system for in-situ CO<sub>2</sub> and CH<sub>4</sub> measurements and calculation method for concentrations and uncertainty.

\*Motoki Sasakawa<sup>1</sup>, Noritsugu Tsuda<sup>2</sup>, Toshinobu Machida<sup>1</sup>, Mikhail Arshinov<sup>3</sup>, Davidov Denis<sup>3</sup>, Alexandrov Fofonov<sup>3</sup> (1.NIES National Institute of Environmental Studies, 2.Global Environmental Forum, 3.Institute of Atmospheric Optics, Russian Academy of Sciences, Siberian Branch, Russia) Keywords:Carbon dioxide, Methane, Siberia, in-situ measurement, CRDS

Continuous measurements of  $CO_2$  and  $CH_4$  concentration have been carried out using a tower network in Siberia (JR-STATION: Japan-Russia Siberian Tall Tower Inland Observation Network) to study the spatial and temporal variations of  $CO_2$  and  $CH_4$  and estimate the distribution of the flux over this vast area (Sasakawa et al., 2010, 2012, 2013) where only a few atmospheric investigations had been made. Transport of working standard gases to remote sites is a significant issue. Thus, to reduce the consumption of the gases, Watai et al. (2010) developed a system that utilizes in-situ air as sub-working standard gas to track the baseline drift of an NDIR. The calibrations using the working standard gases were carried out twice a day. Watai et al. (2010) installed the system at the first tower site in West Siberia (Berezorechka). We modified the working gas saving system and further added a  $CH_4$  sensor (Suto et al., 2010), then expanded the tower network. We evaluated the uncertainty of the concentrations obtained from this modified system. From the year of 2015, we installed a Cavity Ring-Down Spectroscopy (CRDS; Picarro inc.) at Karasevoe, Demyanskoe, and Noyabrsk. We thus validate the recent data with the data by the CRDSs.

## References

Sasakawa, M. et al., Tellus **62B**, 403-416, 2010. Sasakawa, M. et al., Tellus **64B**, doi:10.3402/tellusb.v64i0.17514, 2012. Sasakawa, M. et al., J. Geophys. Res. **118**, 1-10, doi:10.1002/jgrd.50755, 2013. Suto, H. et al., J. Atmos. Ocean. Tech. **27**, 1175-1184, 2010. Watai, T. et al., Atmos. Ocean. Tech. **27**, 843-855, 2010.