

---

[JJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-AS Atmospheric Sciences, Meteorology & Atmospheric Environment

## [A-AS07]Stratosphere-troposphere Processes And their Role in Climate

convener:Shingo Watanabe(Japan Agency for Marine-Earth Science and Technology), Yoshio Kawatani(Japan Agency for Marine-Earth Science and Technology), Takashi Sekiya(国立研究開発法人 海洋研究開発機構, 共同), Kaoru Sato(Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The Stratosphere-troposphere Processes And their Role in Climate (SPARC) is one of the major projects of the World Climate Research Programme (WCRP), and is characterized by its focus on chemical and dynamical coupling in the stratosphere and troposphere. In this session, we welcome presentations on various processes in the stratosphere and troposphere.

---

## [AAS07-P06]Super-pressure balloon observation of gravity waves over the Antarctic

\*Yoshihiro Tomikawa<sup>1,2</sup>, Kaoru Sato<sup>3</sup>, Yoshitaka Saito<sup>4</sup>, Isao Murata<sup>5</sup>, Naohiko Hirasawa<sup>1,2</sup>, Masashi Kohma<sup>3</sup> (1.National Institute of Polar Research, 2.SOKENDAI (The Graduate University for Advanced Studies), 3.The University of Tokyo, 4.JAXA/ISAS, 5.Tohoku University)

Keywords:Super-pressure balloon, gravity wave, Antarctic

Recently super-pressure balloons (SPBs) have been developed, which can float at a constant density surface in the troposphere and stratosphere for long duration (i.e., several months). They can follow Lagrangian motions of air parcels, which is beneficial for gravity wave studies. Gravity wave is one of uncertain factors in current climate models, in which it is required to obtain its stochastic features as well as its spatial and temporal mean behavior. SPBs enable us to obtain stochastic features of gravity waves in a full frequency range from Brunt-Vaisala frequency to inertial frequency. Our group proposes new SPB observations in the Antarctic.