

[JJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-CC Cryospheric Sciences & Cold District Environment

[A-CC29]Ice cores and paleoenvironmental modeling

convener:Ryu Uemura(University of the Ryukyus), Kenji Kawamura(National Institute of Polar Research, Research Organization of Information and Systems), Ayako Abe-Ouchi(東京大学大気海洋研究所, 共同), Nozomu Takeuchi(Chiba University)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Analyses of ice cores from polar and mountain regions have contributed to the reconstruction and understanding of the past environmental changes on timescales from years to several hundred thousand years. In this session, we welcome paleoenvironmental studies using ice cores and paleoclimatic modeling. Studies on reconstruction methods, recording processes and new paleoenvironmental proxies, technical aspects of paleo-modeling are also welcomed. Studies with marine sediment cores, terrestrial sediment cores and tree-rings on similar timescales are also important and welcomed, in order to discuss past environmental changes from multidisciplinary viewpoints.

[ACC29-P11]Iron fertilization and atmospheric CO₂ change during Heinrich event: a model study

*Akitomo Yamamoto¹, Ayako Abe-Ouchi^{2,1} (1.Japan Agency for Marine-Earth Science and TechnologyAtmosphere and Ocean Research Institute, 2.Atmosphere and Ocean Research Institute, The University of Tokyo)

Keywords:ocean carbon cycle, glacial-interglacial cycle, iron cycle, ocean biogeochemical model

Weakening of the Atlantic Meridional Overturning Circulation (AMOC) during Heinrich events was often accompanied by the atmospheric CO₂ increase of 10±20 ppm over 1000 years (Ahn and Brook, 2008). However, previous modeling studies display conflicting atmospheric CO₂ responses to an AMOC shut down. Recent paleoproxy records suggest that dwindling iron fertilization by dust in the Southern Ocean can also explain millennial-scale CO₂ oscillations (Martinez-Garcia et al., 2014). In this study, we investigate the response of atmospheric CO₂ to changes in ocean circulation and dust-borne iron supply using numerical models. In association with the AMOC weakening from 26 Sv to 6 Sv, the atmospheric CO₂ decreases by 0.5 ppmv over 1000 years. Combining AMOC weakening and decrease in dust-borne iron supply, the atmospheric CO₂ increases by 16 ppm over 1000 years. Reduction in export production associated with dwindling iron supply in the Southern Ocean release carbon from the deep sea, contributing to the rise of atmospheric CO₂. Reduction in export production also increases dissolved oxygen in the Antarctic Bottom Water, which is consistent with proxy records. Our results support the contribution of changes in dust-borne iron supply to the millennial CO₂ change in the glacial periods.