
[EE] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-CG Complex & General

[A-CG34] Extratropical oceans and atmosphere

convener: Hatsumi Nishikawa (Institute of Low Temperature Science, Hokkaido University), Yoshi N Sasaki (Hokkaido University), Satoru Okajima (東京大学先端科学技術研究センター, 共同), Thomas Spengler (University of Bergen)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The extratropical ocean had been considered passive to atmospheric variability. Recent studies, however, revealed some active role of the extratropical ocean in modulating the atmosphere. The goal of this session is to deepen our understanding of the air-sea interaction in the extratropics. A wide variety of researches whose topics range from mesoscale to basin-scale, and from daily to global warming are welcomed. Researches on cloud, aerosol, and ecosystem related to the extratropical air-sea interaction are also welcomed.

[ACG34-P06] Mechanisms for Maintaining Basin-Scale Atmospheric Response to Decadal Variability of the North Pacific Oceanic Frontal Zone

*Satoru Okajima¹, Hisashi Nakamura¹, Kazuaki Nishii², Takafumi Miyasaka¹, Akira Kuwano-Yoshida³, Bunmei Taguchi¹, Masato Mori¹, Yu Kosaka¹ (1. Research Center for Advanced Science and Technology, 2. Mie Univ., 3. Kyoto Univ.)

Keywords: Air-Sea Interaction, Oceanic Front

Through the following two ensemble experiments with an atmospheric general circulation model (AGCM), mechanisms for the maintenance of a large-scale wintertime atmospheric response to warm sea surface temperature (SST) anomalies associated with decadal-scale poleward displacement of the North Pacific subarctic frontal zone (SAFZ) has been investigated. The ensemble experiments are as follows: one with climatological-mean SST and the other with positive SST anomalies along the SAFZ prescribed on top of the climatological-mean SST.

As actually observed, the simulated January ensemble response over the North Pacific is anticyclonic throughout the depth of the troposphere, although its amplitude is smaller. This response is maintained through energy conversion from the ensemble climatological-mean circulation realized under the climatological SST as well as feedback from anomalous transient eddy activity, suggesting that the response may have a characteristic as a preferred mode of variability (or "dynamical mode");. Conversions of both available potential energy and kinetic energy from the climatological-mean state are important for the observed anomaly, while the latter is less pronounced for the model response. Net transient feedback forcing is also important for both the observed anomaly and simulated response. These results imply that moderate horizontal resolution (~1 degree) AGCM may be able to simulate a basin-scale atmospheric response to the SAFZ SST anomaly through synoptic and basin-scale dynamical processes. Weaker PNA-like internal variability in the model may lead to the weaker response, suggesting that misrepresentation of intrinsic atmospheric variability can affect the model response to the SST anomaly.