
[EJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-CG Complex & General

[A-CG38]Science in the Arctic Region

convener:Shun Tsutaki(The University of Tokyo), NAOYA KANNA(Arctic Research Center, Hokkaido University), Shunsuke Tei(北海道大学 北極域研究センター, 共同), Tetsu Nakamura(Faculty of Environmental Earth Science, Hokkaido University)

Thu. May 24, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The Arctic and circumpolar region is the key area for the study of global change because the anthropogenic impact is projected to be the largest in this area due to the complicated feedback processes of the nature. A number of international and interdisciplinary research projects have been conducted for the studies on the land-atmosphere-ocean system. In order to understand the feedback processes occurring in the Arctic and to project the global warming in the future, we need to establish the intense observational network and to exchange the knowledge and information by combining the different scientific communities under the common interest of the Arctic. The objectives of this session are 1) to exchange our knowledge on the observational facts and integrated modelling and 2) to deepen our understanding on wide range of natural sciences related to the Arctic and the circumpolar region. Studies on humanities, social sciences, and interdisciplinary fields are also welcomed.

[ACG38-P17]Optical tracer method to distinguish fresh water sources in the Pacific Arctic region

*Amane Fujiwara¹, Atsushi Matsuoka³, Shigeto Nishino¹, Hisatomo Waga², Toru Hirawake², Takashi Kikuchi¹ (1.Japan Agency for Marine-Earth Science and Technology, 2.Faculty of Fisheries Sciences, Hokkaido University, 3.Takuvik Joint International Laboratory, Université Laval)

Keywords:absorption of CDOM, Pacific Arctic region, fresh water sources

Recent freshening of the Arctic Ocean caused notable changes in ocean environment and circulations. Both increases in sea ice melt water and river discharge are known to have a large effect on the freshening. Chemical tracers, oxygen isotope ratio and total alkalinity (TA), are used to distinguish fresh water sources to understand their contribution to the freshening and distribution of the water masses. However, estimation of fresh water sources using chemical tracers has a disadvantage in high frequent sampling, because it takes considerable time to measure. In this study, we propose a method to discriminate water sources using absorption of CDOM (a_{CDOM}) for the Pacific Arctic region. We obtained salinity, TA, and a_{CDOM} at 443 nm ($a_{\text{CDOM}}(443)$) from the seven cruises of R/V Mirai conducted in the Pacific Arctic region during late summer. Fraction of sea ice melt water (f_{SIM}) and other fresh water (f_{OF}) were calculated from their known end-member values of salinity and TA. We identified $a_{\text{CDOM}}(443)$ end-member values of f_{SIM} and f_{OF} , which were unknown, applying a least-square method to the relationship between fresh water fraction and $a_{\text{CDOM}}(443)$. Thus, f_{SIM} and f_{OF} became retrievable from salinity and $a_{\text{CDOM}}(443)$ instead of TA. We also evaluated the accuracy of the optical tracer method comparing fresh water fraction values calculated from chemical and optical tracers, and obtained equivalent results to the chemical tracer method. Estimation of fresh water sources with higher frequent monitoring using a_{CDOM} might advance the comprehension of ocean circulation and related chemical- and biological system in the Arctic.