[JJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-HW Hydrology & Water Environment

[A-HW24]Hydrological change after the 2016 Kumamoto earthquake convener:Jun Shimada(Graduate school of Science and Technology, Kumamoto University), Kei Nakagawa(Graduate School of Fisheries Science and Environmental Studies, Nagasaki University), Takahiro Hosono(熊本大学大学院先導機構, 共同)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) More than two years have been passed after the occurrence of 2016 Kumamoto earthquake. Many investigators have been investigated the cause of observed coseismic hydrological changes such as spring lake dry up, groundwater level drop and rise. We also found groundwater quality changes before and after the quake and this information has been becoming accumulated. In fact, highly densed groundwater monitoring network installed in Kumamoto enables us to grasp comprehensive view of coseismic hydrological responses in very high resolution, so that, the results of these studies have high potential impact to this academic area globally. In this sense, we are welcome for all topics regarding coseismic hydrological changes after or even before the 2016 Kumamoto earthquake from broad point of view including hydrological cycle, deep water and hydrothermal water contribution, subsurface temperature, water quality, isotopes and microbiology. Topics of earthquake prediction and crustal defamation mechanism, surface morphological change in relation to hydrological changes are also welcomed.

[AHW24-P01]Elucidation of groundwater discharge mechanism after the Kumamoto earthquake in northwest Aso

*Koki Okamura¹, Takahiro Hosono² (1.Kumamoto University Graduate School of Science and Technology, 2.Kumamoto University Prioryty Organization for Innovation and Excellence) Keywords:Kumamoto earthquake, northwest Aso, new spring water

The Kumamoto earthquake that occurred on April 14 and 16, 2016, has had major impact on the groundwater environment. The influence of the earthquake was not only on the Kumamoto area, but also the Aso area which is far from the epicenters. It was reported that the flow rate of groundwater increased and some hot spring sources stopped flowing at the time. In the northwestern part of the Aso area, horizontal movement of the topography to the northwest which occurred locally after the main shock was accompanied by compression and extension fields resulting in many cracks and sinks. The appearance of new spring water was confirmed at several places. Previous studies have elucidated the structural changes in the northwestern part of Aso. No hydrological studies on the new spring waters has been carried out yet. This study therefore attempts to elucidate the occurrence of the new springs. General water quality data, isotope analysis, topographical changes and reports on underground structural changes after the earthquake in the northwestern part of Aso are used to decipher the flow and discharge mechanism of the new springs.

The new spring water was classified into two (2); the mountain spring water at the foot of the outer ring mountain, and the plain spring water confirmed at the plain spreading across paddy fields. These waters showed different quality compositions. The mountain spring water was of low concentration Ca- HCO_3 type (the water type of common mountain stream waters), originating from the Somma. On the other hand, the plain spring water showed a high concentration of Ca- SO_4 type originating from the central volcanic hill. It is speculated that this is underground water influenced by volcanic activity. Both waters were of the same quality as spring waters reported so far of the Aso caldera, as well as the distribution area.

Sulfur isotopic ratios were used as tracers to clarify the mechanism for the high concentrated water found only on the plain. The δ³⁴S value varied depending on the site. One point showed relatively high δ³⁴S value (10 ‰ to 17 ‰) and another showed relatively low δ³⁴S value (2.5 ‰). The new spring water in the Kario area has the same isotopic composition as that of hot spring water in the surrounding area, and this may be due to the influence of a heat source similar to that of the hot spring water during the springing process. In a previous study, the δ³⁴S of spring water recharged from the Aso volcano, which is a typical heat source in the Aso caldera, was approximately 5 to 10 ‰. Relatively, spring water in the northwestern part in this study have higher δ³⁴S values, suggesting the possible presence of a heat source different from that of the central crater in the area.

In identifying reasons for the groundwater discharge after the earthquake, different factors were considered for both the mountain spring water and plain spring water based on flow system and water quality characteristics. The release of groundwater reserved beneath the mountain due to seismic motion and the possible formation of " water pathways" from the earthquake could be the reason for the occurrence of the new springs. New spring waters in the paddy fields of the plain whose discharge rate show no decreasing tendency even a year after the earthquake, taking into consideration seasonal variation, indicates that major underground structural changes have occurred. It can be inferred that new discharge route for groundwater has been created.