

---

[JJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-HW Hydrology & Water Environment

## [A-HW26]Water Environment and Geology in Urban Areas

convener:Takeshi Hayashi(Faculty of Education and Human Studies, Akita University), Kei Nishida(Interdisciplinary Centre for River Basin Environment, Interdisciplinary Graduate School, University of Yamanashi), Hiroaki SUZUKI(日本工営株式会社 中央研究所, 共同)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The scope of this session is to create an interdisciplinary forum on the most recent advances in water environment and environmental geology research in urban areas. Various kinds of studies concerning environmental issues on water and geology in urban areas (e.g. water balance, water cycle, water resource development and management, inundations, hydrogeology, pollution and remediation, geohazard, basic law on the water cycle) are welcome from academia, industry, and government as well as wider geographic diversity.

---

## [AHW26-P04]Salinity of pore water in low permeable alluvial sediments in the inner Tokyo Bay

\*Takeshi Hayashi<sup>1</sup>, Akinobu Miyakoshi<sup>2</sup>, Yoshinori MIYACHI<sup>2</sup>, Junko Komatsubara<sup>2</sup>, Masaya Yasuhara<sup>3</sup>  
(1.Faculty of Education and Human Studies, Akita University, 2.GSJ, AIST, 3.Faculty of Geo-environmental Science, Rissho University)

Keywords:Tokyo Bay area, Alluvium, pore water, salinity

Geochemical characteristics of pore water in sediments including alluvium reflect not only sedimentary environment but also various processes such as diagenesis, advection and dispersion due to groundwater flow, geochemical interactions with sediments, and diffusion. The changes of geochemical characteristics of pore water have been widely observed not only in “aquifer (e.g. sand and gravel)” but also in “aquitard (e.g. silt and clay)” around the world. In addition, in coastal urban areas, human activities such as reclamation and groundwater development have complicated the geochemical processes. However, the complicated processes in the urban area have not been clarified. Geological Survey of Japan, AIST has carried out all-core boring in the inner Tokyo Bay area (from Tokyo lowland to Chiba city) to grasp lithology, sedimentary environment and geological structure. We also have focused on geochemistry of pore water in the alluvium in this area, to understand processes and mechanisms of solute transport and geochemical reactions. We will present the results of estimation of salinity distribution and the mechanisms of it in the alluvium of this area.

As for core samples that were obtained in Chiba area (Funabashi city and Chiba city), Electric Conductivity (EC) of pore water in low permeable alluvial sediments (silt and clay) was from 65 to over 2,000 mS/m. Over 2,000 mS/m means that EC exceeded the measuring limit of the instrument. These values were higher than that of suspensions of core samples obtained at the Tokyo lowland (67 to 1,000 mS/m; Uchida et al., 2011). Salinity was also considered to be high in the upper part of alluvium. These results suggest that desalination has occurred in the alluvium of Chiba area.