
[JJ] Evening Poster | A (Atmospheric and Hydrospheric Sciences) | A-OS Ocean Sciences & Ocean Environment

[A-OS17] Ocean circulation and material cycle in coastal seas

convener: Akihiko Morimoto (Ehime University), Kiyoshi Tanaka (University of Tokyo), Yuichi Hayami (佐賀大学, 共同), Kazuhiko Ichimi (Faculty of Agriculture, Kagawa University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Costal seas have high biodiversity and high productivity, while various environmental problems such as eutrophication, red tide, and hypoxia are occurred by human activity. In order to understand material cycle and various phenomena and to maintain sustainable environment in costal seas, interdisciplinary research is needed. In the present session, we invite researchers who are interested in physical phenomena, material cycle, biological responsive dynamics, and those interactions in coastal seas. We do not confine study field, area, and method and welcome presentations of wide-range topics. We try to commoditize knowledge in terms of phenomena in coastal seas based on the presentations and discussion.

[AOS17-P03] Hydrographic observation on the shelf off Otsuchi Bay in northeastern Japan

*Hajime Nishigaki¹, Yuta Takada², Kiyoshi Tanaka³, Daigo Yanagimoto³, Shinzo Fujio³, Kei Nishina³

(1. Faculty of Science and Technology, Oita University, 2. Faculty of Education and Welfare Science, Oita University, 3. Atmosphere and Ocean Research Institute, The University of Tokyo)

Keywords: baroclinic tide, shelf, Sanriku coast

The Sanriku ria coast is located on the eastern coast of northeastern Japan facing North Pacific. Fisheries such as farming of seaweeds and shellfish are active in this area. Therefore, the topic of water circulation and exchange is important for the fishery and recovery from the disaster of the 2011 tsunamis, in addition to scientific interests. Otsuchi Bay is one of the bays in this area at about 39°20' N, whose size is about 8 km in east-west and 3 km in north-south and up to 70 m in depth. This study aims to reveal water circulation in the shelf area off Otsuchi Bay by a hydrographic observation. Within Otsuchi Bay, baroclinic circulations are revealed by observations (e.g. Otake et al., 2009; Tanaka et al., 2017; Ishizu et al., 2017). In summer, baroclinic tidal currents are observed with offshore/inshore upper current and inshore/offshore lower current. Together with a thin surface layer of a few meters, it makes a three-layer structure (Tanaka et al., 2017). However, it is not clear how this current reaches the mouth of the bay because observations in the offshore shelf area is inadequate. This study makes the three-dimensional observation with finest resolution ever in a shelf area in the Sanriku ria coast.

Hydrographic observation was done with R/V Yayoi in the area of 4.3 km in east-west and 5.6 km in north-south, where the depth is between 50 m and 125 m. It was on July 26, 2017 in the morning (8:11-11:00) and the afternoon (1:12-2:13). In the morning, three-dimensional CTD observation was made. In the afternoon, observation is limited on the coastal meridional line with a finer horizontal resolution. Measurements of the velocity by a shipboard ADCP (300 kHz) are also made during the cruise. A nearby location, Kamaishi had a low tide at about 11:30 am with an intermediate tidal range. So, the morning/afternoon cruise corresponds to ebb/flood tide at the coast near the site.

Observed distribution of the salinity and temperature has two haloclines with salinity of about 33.60 and 33.70, respectively. The former lies at the depth of a few meters. The depth of the latter varies from 20 m to 60 m. The salinity and temperature of the lower halocline and the water below is categorized in the Tsugaru Warm Current water system of Hanawa and Mitsudera (1987). Distributions of the zonal velocity by the shipboard ADCP shows baroclinic structure with upper offshore flow and

lower inshore flow. The boundaries between them correspond to the lower halocline in CTD observations. These results are consistent with the three-layer structure in inshore Otsuchi Bay revealed by Tanaka et al. (2017). The morning and afternoon observations of the coastal meridional line show significant salinity increase and temperature decrease in the deeper portion, especially in its northern part. The morning observations in the offshore area suggest that the water with high salinity and low temperature came from offshore region with upslope movement.

References.

Hanawa and Mitsudera, 1987, JOSJ, 42.

Ishizu et al., 2017, JO, 73.

Otobe et al., 2009, Coast Mar Sci, 33.

Tanaka et al., 2017, JO, 73.