
[EE] Evening Poster | B (Biogeosciences) | B-AO Astrobiology & the Origin of Life

[B-AO01]Astrobiology

convener:Hikaru Yabuta(Hiroshima University, Department of Earth and Planetary Systems Science), Seiji Sugita(Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo), Misato Fukagawa(名古屋大学, 共同), Fujishima Kosuke(Tokyo Institute of Technology, Earth-Life Science Institute)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Twenty years have passed since when the field of Astrobiology, which aims to unveil the origins, evolution, and habitability of life by integrating multidisciplinary fields, was established. Origins of Life are currently being re-conceptualized via expansion of prebiotic chemistry to systems chemistry and chemical space. Besides their relationship to life's building blocks, it is expected to demonstrate the significant roles of organic molecules in the history of planetary formation. The linkages among the variations in chemical compositions of deep-sea hydrothermal environments, geological settings, and ecological systems were systematically investigated. Cassini, which accomplished in the long-term explorations of the planets bearing liquid, had "Grand Finale" this year. Discoveries of extrasolar planets have been dramatically increased to date.

Originally, Astrobiology does not need a specific science category. We therefore aim to make this session so that Earth and Planetary scientists from all the categories join for discussing 'where we came from and where we are going' and for making novel integrated researches.

For the next stage of Astrobiology, presentations on the instrument development in space explorations, comparative studies of solar system and exoplanets, etc, will be very much welcome.

[BAO01-P09]Experimental study on role of sulfide-peptide complexes in the first metabolism

*Kensuke Igarashi¹ (1.National Institute of Advanced Industrial Science & Technology)

Keywords:Origin of life, Iron sulfide, Primitive enzyme, Iron-sulfur protein, Chemical evolution, Acetyl-CoA pathway

Metal sulfides are thought to have played an essential role in chemical evolution processes leading to the origin of life (the iron-sulfur world hypothesis). Greigite (Fe_3S_4) contains the cubic Fe_4S_4 unit, which is similar to the iron sulfur cluster, which constitutes the catalytic center of various iron-sulfur proteins such as carbon monoxide dehydrogenase/acetyl-CoA synthase. Although greigite is thought to have been the origin of iron-sulfur clusters, the catalytic activity of this mineral has not been assayed for hydrothermal reactions in which inorganic carbons (i.e., CO_2 and CO) are reduced to form organics. In chemical evolution processes, it has been hypothesized that metal sulfides were conjugated with peptides to form sulfides-peptide complexes, which had greater stability and higher catalytic activity than the metal sulfides themselves. However, no experimental evidence that supports this hypothesis has been reported. Thus, the aims of this study are to (i) obtain a more comprehensive understanding of the catalytic nature of metal sulfides, including greigite, under hydrothermal conditions wherein inorganic carbons are converted to organics and (ii) obtain evidence for enhancement of the catalytic activity of metal sulfides by complex formation with peptides.

The catalytic activity of metal sulfides (FeS , Fe_3S_4 , NiFe_2S_4 , FeS_2 , CoS , NiS , and ZnS) for the conversion of CO_2 or CO was assayed under various hydrothermal conditions (20–100°C, pH 4–10, 2–100 atm, with/without reductants). Metal sulfides containing Fe and/or Ni catalyzed CO_2 reduction by H_2 to afford formate. Among them, the greigite species (Fe_3S_4 and NiFe_2S_4) showed the

highest catalytic activity. Similarly, greigite catalyzed CO reduction by HS⁻ more efficiently than did any other metal sulfides tested, producing 2–6 mM acetate after 24 h reaction under 100°C and 2 atm. At higher reaction pressures, CO₂ was predominantly reduced by H₂ to methane, indicating dissolved gas concentration strongly influence reaction products. These results suggest that greigite has superior catalytic activity and plays a central role in establishing the first metabolism. In order to construct sulfide-peptide complexes, greigite-binding peptides were screened by using phage display random peptide libraries. After three rounds of biopanning, several consensus peptide sequences that were rich in polar amino acids (especially histidine) were obtained and found to attach to greigite. Preliminary experiments to study sulfide-peptide complexes are underway, and the results would be reported in due course.