[EE] Evening Poster | B (Biogeosciences) | B-CG Complex & General

## [B-CG07]Earth and Planetary Science Frontiers for Life and Global Environment

convener:Yoshinori Takano(Japan Agency for Marine-Earth Science and Technology (JAMSTEC)), Yohey Suzuki(Graduate School of Science, The University of Tokyo), Keisuke Fukushi(金沢大学環日本海域環境研究センター, 共同), Shingo Kato(RIKEN)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) Earth and planetary science is currently capable of sampling from three frontiers: space, deep ocean and deep underground for the study of the origin and evolution of life. Next generation sequencing technology of nucleic acids and nanotechnology-driven innovation of solid characterizations are very powerful to expand our understanding of many fields of earth and planetary sciences.

It is also important for us to be aware of global warming and climate change that dramatically affect biodiversity, biological activities and elemental cycling.

The aim of this session is to cross the border of science sections to explore the frontiers of earth and planetary science.

## [BCG07-P02]Reconstructing Ecological Responses to Last Global Warming Recorded in Japan Sea Sediment through Ancient DNA Analysis

\*Mariko Kouduka<sup>1</sup>, Yohey Suzuki<sup>1</sup> (1.Department of Earth and Planetary Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, Japan.)

Keywords: Japan sea, Ancient DNA, Last Global Warming, anoxic bottom water

The advancements of scientific ocean drilling towards climate and ocean changes through sedimentological, paleontological and biogeochemical approaches and biosphere frontiers through nextgeneration sequencing of genomic DNA from the deep biosphere lead to an emerging field of ancient DNA genomics. In semi-closed ocean basins, climate-driven changes in marine ecosystem are pronounced because of the channel closure and physicochemical stratification based on salinity and temperature. The lack of O<sub>2</sub> to benthic organisms results in the formation of unturbated sediment with laminated layers, each of which tends to record the past ocean change at high temporal resolution. As pilot studies demonstrated that DNA and other biomolecules are well preserved at geological time scale under O<sub>2</sub>deprived conditions, ancient DNA genomics is becoming a new powerful tool to decode the past ecological change. We investigated ancient DNA obtained from thinly laminated layers deposited in the Japan Sea across the most recent global warming event rapidly occurred after Younger Dryas. Metabarcoding of 18S rRNA gene sequences clarified that across the transition from oxygenated to anoxic bottom water conditions, dominant protist populations were shifted from radiolarians (Acantharia) to diatoms (Thalassiosirales) and unclassified Stramenopiles. These results suggest that the inflow of Tsushima Current with nutrient-rich, warm seawater and rapid global warming triggered the complex anoxiaecology interactions. It is also suggested that the long-term ecological change reconstructed in this study provides realistic information to predict ecosystem perturbation caused by current global warming.