
[JJ] Evening Poster | B (Biogeosciences) | B-CG Complex & General

[B-CG09]Decoding the history of Earth: From Hadean to the present

convener: Tsuyoshi Komiya (Department of Earth Science & Astronomy Graduate School of Arts and Sciences The University of Tokyo), Yasuhiro Kato (Department of Systems Innovation, Graduate School of Engineering, University of Tokyo), Katsuhiko Suzuki (国立研究開発法人海洋研究開発機構・海底資源研究開発センター)

Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The latest results of Earth's evolution and geological processes through 4.6 billion years from Hadean to Modern, based on various approaches including fieldworks, chemical analyses, experiments and computer simulation, will be presented. In this session, we aim to discuss and understand causal relationships and interplay among the evolution of Earth's deep interior, changes in the surface environments, and development and evolution of life. Wide-ranging topics are accepted.

[BCG09-P14]Geochemical characterization of modern black mud by Independent Component Analysis

*Moei Yano¹, Kazutaka Yasukawa^{1,2}, Koichiro Fujinaga^{2,1}, Kentaro Nakamura¹, Minoru Ikehara³, Yasuhiro Kato^{1,2} (1. Department of Systems Innovation, School of Engineering, The University of Tokyo, 2. Ocean Resources Research Center for Next Generation, Chiba Institute of Technology, 3. Center for Advanced Marine Core Research, Kochi University)

Keywords: redox sensitive metals, black mud, black shale, Independent Component Analysis

Black shales are enriched in some kinds of redox sensitive metals as well as organic carbon and sulfides, and have been formed through the Earth history with the emergence of oxygen-depleted environment. The concentration of these metals has been attributed to an adsorption by organic matter, incorporation to or formation of sulfide minerals, or a precipitation as insoluble compounds such as oxyhydroxides. In the modern ocean, black mud that is enriched in organic carbon and/or sulfides deposits in certain sea areas associated with an oxygen-depleted water mass, such as a silled basin or an upwelling area. These black muds likely constitute comparable counterparts of the ancient black shales. Therefore, to characterize the geochemical features of these black muds may provide a new insight into the past oceanic environmental changes.

In this study, we determined contents of major and trace elements, total organic carbon (TOC), and total sulfur (TS) of black mud and shale samples collected from a variety of geographic conditions and redox states in the water mass. In order to characterize the geochemistry of black mud and shale, we also applied Independent Component Analysis to the multi-elemental data set. We will present the analytical result and discuss its implications for the oceanic environmental changes.