

---

 [JJ] Evening Poster | B (Biogeosciences) | B-PT Paleontology

## [B-PT06]Biotic History

convener: Isao Motoyama (Department of Earth and Environmental Sciences, Yamagata University), Takao Ubukata (Division of Geology & Mineralogy, Department of Earth & Planetary Sciences, Kyoto University), Kazuyoshi Moriya (早稲田大学 教育・総合科学学術院 地球科学専修)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The Biotic History session covers all aspects of ancient life and the history of biosphere through the geologic time. The study of ancient life is essential for unveiling mysteries of our planet earth. It also provides evidence for evolution of oceans, continents and island arcs. Modern progress in this field has been enhanced by interdisciplinary collaboration with allied sciences, such as paleoceanography and evolutionary biology. Our session intends to be a hub of communication amongst all earth scientists studying the biosphere; we welcome biological and biogeochemical approaches toward the understanding of the history of life.

---

## [BPT06-P05] Reconstruction of pelagic reef biota of the Carboniferous Omi Limestone, Niigata Prefecture, central Japan

\*Yui Takahashi<sup>1,2</sup>, Katsuo Sashida<sup>1</sup>, Sachiko Agematsu<sup>1</sup> (1.University of Tsukuba, 2.Muroto Geopark Promotion committee)

Keywords: Microfossil, Akiyoshi Terrane, Omi Limestone, Paleozoic Era, Carboniferous

Reefs are most biodiversified ecosystems of the modern ocean and their intricate three-dimensional landscapes promote elaborate adaptations of marine animals and complex interactions of them. However, the reefs are the most threatened ecosystems in the world, and the transitions of the reefs from geological time up to the present are one of the major interests of reef studies. Paleozoic pelagic reefs are distributed as reef carbonates in Japanese Islands. We study the Carboniferous Omi Limestone belonging to the Akiyoshi Terrane as one representative here. The Omi Limestone occupies the eastern end of the Akiyoshi Terrane is located in the Itoigawa City, Niigata Prefecture, central Japan. Due to containing rich fossils, the Omi Limestone has been subject to paleontological studies since in the early 1900s and the limestone is a historic place of the Japanese paleontology. The oldest work conducted in 1918 by Hayasaka reported some Carboniferous brachiopod fossils and revealed the presence of the Carboniferous strata in Japan for the first time. This limestone possesses abundant easily-recognizable fossils, but remnants in fossil biota are less distinctive. For that reason, the primary reports themselves of microfossils are few except for ones with biostratigraphical importances such as smaller foraminifers and fusulinaceans. This study aims to reveal the microfossil fauna and reconstruction of the Carboniferous pelagic reef ecosystem based on the integration of micro and known macrofossils. Obtained microfossils fauna is composed of following taxa: conodonts, ostracods, actinopterygians and chondrichthyans, mollusks, echinoids, holothuroids, ophiuroids, sponge spicules and chitinozoas. Classes Ostracoda, Actinopterygii, Chondrichthys, Echinoidea, Holothuroidea, Ophiuroidea and Chitinozoa are first report from the limestone. Based on obtained conodonts, four middle Carboniferous conodont biozones are established. We reconstruct the paleoecosystem of the middle Carboniferous Omi Limestone from the perspectives of paleoecology of fossil taxa. We also investigate the paleogeography of the middle Carboniferous Omi Limestone based on the regional and international correlations of benthic faunas such as holothuroid sclerites and ostaracods.