[JJ] Evening Poster | B (Biogeosciences) | B-PT Paleontology

[B-PT06]Biotic History

convener:Isao Motoyama(Department of Earth and Environmental Sciences, Yamagata University), Takao Ubukata(Division of Geology & Mineralogy, Department of Earth & Planetary Sciences, Kyoto University), Kazuyoshi Moriya(早稲田大学 教育・総合科学学術院 地球科学専修)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe) The Biotic History session covers all aspects of ancient life and the history of biosphere through the geologic time. The study of ancient life is essential for unveiling mysteries of our planet earth. It also provides evidence for evolution of oceans, continents and island arcs. Modern progress in this field has been enhanced by interdisciplinary collaboration with allied sciences, such as paleoceanography and evolutionary biology. Our session intends to be a hub of communication amongst all earth scientists studying the biosphere; we welcome biological and biogeochemical approaches toward the understanding of the history of life.

[BPT06-P07]Radio- and stable carbon isotopic responses in experimentally-cultured bivalves for the understanding of acidification effect on bivalve nutrient uptake and biomineralization

*Kozue Nishida^{1,2}, Yue Chin Chew³, Yusuke Yokoyama³, Atsushi Suzuki⁴, Yosuke Miyairi³, Shoko Hirabayashi⁷, Masahiro Hayashi⁵, Yukihiro Nojiri⁶ (1.National Institute of Technology, Ibaraki College, 2.JSPS, 3.AORI, UTokyo, 4.AIST, 5.MERI, 6.Hirosaki Univ., 7.UTokyo) Keywords:Mollusca, Radio isotope, Stable carbon isotope

The effects of ocean acidification on bivalve growth and calcification have been widely debated in recent years due to potential impacts on the aquaculture sector and thus marine biodiversity. This study aims to elucidate the effects of ocean acidification on bivalve nutrient assimilation into its various components via laboratory culture experiments at six different pCO_2 levels at 25 °C (332, 463, 653, 872, 1137, and 1337 μatm). Radio carbon (Δ¹⁴C) and stable carbon (δ¹³C) isotopic analysis will be undertaken to investigate: (1) the proportion of incorporation of carbon from ambient DIC and metabolic CO_2 into bivalve shell and tissues; and (2) effects of changing pCO_2 concentrations on carbon incorporation. Our research marks the first attempt to trace partitioning of nutrients within a single species via application of bomb-pulse radiocarbon principles. By using modern samples in a culture experiment of ocean acidification by using CO_2 gas derived from fossil fuels, differences in contribution from end-members of DIC and metabolic CO_2 can be clearly resolved in high resolution. Combination of radiocarbon techniques with stable isotopic analysis will allow biological contributions to be parsed from a geochemical perspective, providing a comprehensive understanding of bivalve nutrient assimilation.

Specimens of the filter-feeding clam *Scapharca broughtonii* (Bivalvia: Arcidae) were cultured in aquaria for 8 weeks, with a novel high-precision *p*CO₂ control system which maintained steady CO₂ levels. Bivalve shells and soft tissues, as well as the relevant water and plankton feed samples, were then analysed for δ¹³C in an Isotope Ratio Mass Spectrometer (IRMS) and Δ¹⁴C in an Accelerator Mass Spectrometer (AMS) to examine variations in end-member contributions to the bivalve components.

Shell carbon was found to be principally derived from seawater DIC in all pCO₂ conditions. Mantle and

soft tissue isotopic signatures stayed constant across *p*CO₂ concentrations and were primarily correlated with the organic carbon signal. A high degree of correlation between Δ¹⁴C values of bivalve shell and seawater DIC may point to *S. broughtonii*'s suitability as a geochemical proxy of DIC palaeo-concentrations. Using Δ¹⁴C in concert with δ¹³C allowed separation of isotopic fractionations deriving from effects other than those caused by actual changes in end-member contributions. Shell δ¹³C thus might indicate biologically-related fractionation due to physiological adaptations to cope with effects of changes in ambient pH as these dictated the magnitude of this species-specific effect.