
[EE] Evening Poster | H (Human Geosciences) | H-CG Complex & General

[H-CG25]Deltas and estuaries: multidisciplinary analyses of complex river-mouth systems

convener:Yoshiki Saito(Estuary Research Center, Shimane University), Kazuaki Hori(Department of Geography, Graduate School of Environmental Studies, Nagoya University), Guan-Hong Lee(共同), Qing He(State Key Laboratory of Estuarine and Coastal Research, East China Normal University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This session has foci on morphodynamics, material cycling, and sustainability for deltas and estuaries. We invites contributions that advance research on deltas and estuaries as complex river-mouth systems from modern and ancient examples, and develop integrated frameworks for delta & estuary dynamics modeling on various temporal and spatial scales from studies of coastal evolution over the Quaternary to small-scale sediment/material transport processes and also studies based on field observations, numerical simulation and flume studies, and also contributions that promote data collection and sharing for advancing science and local solutions, consider policy and governance issues linked to the sustainable development of deltas and estuaries, and use in-situ and satellite data for guiding modeling and risk assessment.

[HCG25-P08]Human-induced changes in recent sedimentation rates in Bohai Bay, China: implications for coastal development

*Fu Wang¹, Zhu Li Tian¹, Yu Xing Jiang¹, Marshall William², Hong Wang² (1.Tianjin Center of Geological Survey, China Geological Survey, China, 2.Plymouth University Plymouth, United Kingdom)

Keywords:210Pb and 137Cs, sedimentation rate, reclamation

In many countries, coastal planners strive to balance demands among civil, commercial strategy and environmental conversation interests for future development, particularly given the sea level rise in the 21st century. Achieving a sustainable balance is often a dilemma, especially in the low-lying coastal areas, where a large amount of fluvial sediment has been taken away by dams in the river basin. In recent years, we investigated the shore of Bohai Bay in northern China and found large-scale coastal reclamation by the sea, with a severe increase in sea level. To investigate this trend, we obtained sediment cores from near-shore in Bohai Bay, which were dated by ¹³⁷Cs and ²¹⁰Pb radionuclides to determine the sedimentation rates for the last 50 years. The average sedimentation rates of Bohai Bay exceeded 10 mm/y before 1963, which was much higher than the rate of local sea-level rise. However, our results showed an overall decreasing sedimentation rate after 1963, which was not able to compensate for the increasing relative sea-level rise in that time period. In addition, our results revealed that erosion occurred after the 1980s in the shallow sea area of Bohai Bay. We suggest that this situation places the Bohai Bay coast at a much greater risk of inundation and erosion within the next few decades than previously thought, especially in the large new reclamation area. This study may be a case study for many other shallow sea areas of the muddy coast if the sea level continues to rise rapidly and the sediment delivered by rivers continues to decrease.