
 [EE] Evening Poster | H (Human Geosciences) | H-CG Complex & General

[H-CG25]Deltas and estuaries: multidisciplinary analyses of complex river-mouth systems

convener:Yoshiki Saito(Estuary Research Center, Shimane University), Kazuaki Hori(Department of Geography, Graduate School of Environmental Studies, Nagoya University), Guan-Hong Lee(共同), Qing He(State Key Laboratory of Estuarine and Coastal Research, East China Normal University)

Mon. May 21, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

This session has foci on morphodynamics, material cycling, and sustainability for deltas and estuaries. We invites contributions that advance research on deltas and estuaries as complex river-mouth systems from modern and ancient examples, and develop integrated frameworks for delta & estuary dynamics modeling on various temporal and spatial scales from studies of coastal evolution over the Quaternary to small-scale sediment/material transport processes and also studies based on field observations, numerical simulation and flume studies, and also contributions that promote data collection and sharing for advancing science and local solutions, consider policy and governance issues linked to the sustainable development of deltas and estuaries, and use in-situ and satellite data for guiding modeling and risk assessment.

[HCG25-P09]Sediment budgets of the Yellow River delta during 1976–2012 with morphological changes and sediment accumulation rates

Liangyong Zhou^{2,3}, Jian Liu^{2,3}, *Yoshiki Saito¹, Shaobo Diao^{2,3}, Maosheng Gao^{2,3}, Jiandong Qiu^{2,3}, Congliang Xu⁴, Lelong He^{2,3} (1.Estuary Research Center, Shimane University, 2.Key Laboratory of Coastal Wetland Biogeosciences, 3.Qingdao Institute of Marine Geology, 4.Institute of Yellow River Estuary and Coast Science)

Keywords:Topographic changes, ¹³⁷Cs dating, Yellow River delta

The Yellow River (Huanghe) is a major sediment supplier to the Bohai Sea and the Yellow Sea, resulting in the formation of the huge, dynamic delta complex. In 2012 and 2013, extensive bathymetric and high resolution seismic profiles, and sediment cores and surface sediment samples were collected off the new delta lobe in the south Bohai Sea. This study examined the delta sedimentation and morphology of the new lobe and provided a quantitative assessment of the modern fluvial-derived sediment dispersal with mainly core data and bathymetric mapping.

Both the water depth changes and the ages of marine sediments give valuable information on sediment dynamics and sedimentary processes on subaqueous deltas and their adjacent shelf in the Bohai Sea. The two methods were used to estimate the sediment budget in the Yellow River delta shelf during 1976–2012. Since 1976 the river's channel has been located on the east side of delta complex and has built out broad sedimentary lobe. In 2012, extensive bathymetric and high resolution seismic profiles, vibrocores in the survey lines and surface sediments were collected. This study examined the sedimentation and morphology in the modern Yellow River delta and in Laizhou Bay, based on analyses of (1) radionuclides (¹³⁷Cs, ²¹⁰Pb, ¹³⁴Cs), (2) sediment structure and texture, (3) surface sediment distribution pattern, as well as (4) the morphological change during 1976 to 2012. Bathymetric profiles, especially the South-North profiles, revealed the present morphology of the delta front which exceeds previous estimated boundary, and this also was validated on basis of analysis of ¹³⁷Cs in cores.

The ^{137}Cs onset depths corresponding to the depths of lithological changes and morphological changes indicate that it can be a proxy to track the dispersal of Yellow River-derived sediments in the study area. Synthesis of bathymetry, seismic profiles, ^{137}Cs profiles and surface sediment pattern show that the depocenter occurs in the south flank of the Yellow River delta (morphologically a spit) in west of Laizhou Bay, probably resulting from the headland eddy in this area. ^{137}Cs profiles and morphology changes show clearly the distribution of sediment thickness in the whole study area and sediment dispersal pattern from the delta front to the Laizhou Bay since 1976, and ^{210}Pb profiles provide reliable accumulation rates only in shelf area.