

[EE] Evening Poster | H (Human Geosciences) | H-DS Disaster geosciences

[H-DS07] Landslides and related phenomena

convener: Masahiro Chigira (Disaster Prevention Research Institute, Kyoto University), Gonghui Wang (Disaster Prevention Research Institute, Kyoto University), Fumitoshi Imaizumi (静岡大学農学部)
 Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)
 Mass movements, such as landslides, rockfalls, and debris flows, have been occurring extensively in a large number of countries, causing heavy damage. In order to understand them and mitigate induced disasters, we would like to discuss on various issues. We invite contributions that report and discuss on mass movements and related phenomena, focussing on improved understanding of their characteristics; new insights into landslide mechanisms; the development of new approaches to monitoring; novel approaches to behaviour forecasting and prediction; studies of successful landslide management; and the development of methods for hazard and risk evaluation.

[HDS07-P11] Deciphering recent landslide dynamics in the Shirakami Mountains, a World Natural Heritage site, Japan

*Ching-Ying Tsou¹, Taiki Higuchi², Daisuke Higaki¹ (1. Faculty of Agriculture and Life Science, Hirosaki University, 2. NIPPON KOEI CO., LTD)

Keywords: Landslide dynamics, Shirakami Mountains, UAV, SfM, DEM, River Erosion

Larger and deeper landslides are common phenomena, but measurements relevant to landslide dynamics are scarce in the Shirakami Mountains, a World Natural Heritage site, Japan. This study documents recent landslide occurrences (ca. 10 years) of three landslides align on the terrace scarp (at right bank) and have occurred at the outside bend of the meandering channel of the northward-flowing Okawa River. We construct digital surface model (DSM) and orthomosaics at 6.5 cm resolution from UAV imagery obtained on 6 November 2017 using structure-from-motion (SfM) photogrammetric procedures. Then, the DSM was resampled to a 1-m resolution and was used to compare to an airborne LiDAR-derived digital elevation model (DEM) of 1-m resolution acquired on 22 October 2008 to constrain the landslide magnitude and direction of the displacement vectors using CIAS (the Correlation Image Analysis Software; Küb and Vollmer, 2000). Results were also compared with longitudinal profiles collected by a hand-held laser-ranger finder. The computed horizontal displacements range from 0.1 to 9.7 m with a standard deviation of 1.81 m. At the landslide body of the northern landslide the computed vectors suggest a southward movement, however, the explanation for this movement is currently unknown. The longitudinal displacement for a flat, terrace-like topography of 6×7 m within the landslide body of the southern landslide was computed of ~7 m that is appropriately represented as compared with the field measurement. Active displacements were generally constrained at the secondary and side scarps (3–9 m) and the edges of toes (2–8 m), where the retreating movements are well identified by the algorithm. The retreat of landslide toes was likely triggered by lateral river erosion based on the interpretation of changes of river courses by aerial photos and an outcrop exposed on the landslide toe that consist of imbricated, horizontally bedded gravels. Our results indicate that a combination of UAV-based SfM data and available DEM and the image correlation algorithm is useful and effective to understand the landslide dynamics.

This study was funded by the JSPS KAKENHI (Grant Numbers 16K20893). The Lidar DEM used in this study was provided by Geospatial Information Authority of Japan.