
[JJ] Evening Poster | H (Human Geosciences) | H-QR Quaternary research

[H-QR04]Quaternary, Diachronic dynamics of human-environment interactions

convener:Mamoru Koarai(Earth Science course, College of Science, Ibaraki University), Toshihiko Sugai(Department of Natural Environmental Studies, Institute of Environmental Studies, Graduate School of Frontier Science, The University of Tokyo), Kiyohide Mizuno(国立研究開発法人産業技術総合研究所地質情報研究部門, 共同), Minoru YONEDA(The University Museum, The University of Tokyo)
 Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)
 Humans have attained their specific development by indigenous cultures and evolved through environmental adaptation. The session raises issues of human-environmental interactions, views from diverse changes of climate, ocean, land and biota having made striking influence on humans. It welcomes various fields from human-environment change and their chronometric dating among Quaternary disciplines.

[HQR04-P08]Active tectonics and evolutionary processes of the eastern part of Oiso Hills inferred by depositional environments of gravel layers, central Japan

*Masahiro Muraki¹, Taro Funatsu¹, Toshihiko Sugai¹ (1.Univ of Tokyo.)

Keywords:gravel layer, depositional environments, active tectonics, Oiso Hills

Oiso Hills is an uplift block bounded by active faults and uplifted rapidly in the last Quaternary period (Active Fault Research Association, 1991) . In the eastern part of Oiso Hills, there are some active faults. Their continuity is not clear enough because of their complex geological structures (Kaneke. 1971; Mizuno. 2016). We tried to clarify the active tectonics in this area based on the depositional environments of gravel layers and the paleo-geography, reconstructed by analysis of the Pleistocene Kissawa Formation (KF). Firstly, we examined the present Kaname riverbed gravels and Sagami Bay beach gravels. Secondly, we analyzed gravels in KF. Finally we tried to estimate the evolutionary processes of the eastern part of Oiso Hills.

We established a frame of 1m×1m at each gravel sampling site along Kaname River, and the Sagami bay beach, and collected 100 gravels. We also collected 100 gravels from an outcrop within the frame of 1m×1m. Sampling sites were shown in figure 1. We measured the diameter, roundness, degree of weathering, sphericity, and lithology to estimate transportation processes (Sneed · Folk,1958).

The gravels of Kissawa Formation showed the, average diameter, roundness, sphericity of about 2cm, 0.6, 0.7, respectively.- a ratio of mudstone was high subsequently a sandstone, a chart excelled. Average diameter of the Kaname River gravels decreased downstream from around 8cm to around 3cm, and average roundness increased from 0.3 to 0.6. Average sphericity was around 0.7. In gravel species, pyroclastic rocks occupied the majority upstream, and the ratio of sandstone, mudstone, and conglomerate increased in the downstream.

The average diameter of the Sagami Bay beach gravels was around 2.5cm, average roundness was around 0.7, and average sphericity was 0.6, only in the west of the eastern part of Oiso Hills was around 0.7. In gravel species, a sandstone and mudstone excelled, and ratios of mudstone increased towards the west from the Kaname River mouth.

Because mudstone gravel is dominant in KF, and because the average diameter of the gravels of K-F is almost the same as that of the beach gravels, it is more likely to be marine.

Reference: Kaneko S. (1971) Journal of The Geological Society of Japan, Vol. 77, No. 6, p. 345-358.
Active Fault Research Association (1991) University of Tokyo Press, p. 4. Machida · Moriyama (1968)
vol. 41, No. 4, p. 241-257. Mizuno (2016) No. 71, p. 153-160. Sneed ED · Folk RL (1958) Journal of
Geology, Vol. 66, No. 2, p. 114-150.