

[EJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-AG Applied Geosciences

## [M-AG33]Dynamics of radionuclides emitted from Fukushima Dai-ichi Nuclear Power Plant in the environment

convener:Kazuyuki Kita(Faculty of Science, Ibaraki University), Yuichi Onda(Center for Research on Isotopes and Environmental Dynamics, University of Tsukuba), SHINOHARA ATSUSHI(大阪大学, 共同), Daisuke Tsumune(Central Research Institute of Electric Power Industry)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The Great East Japan Earthquake caused the severe accident in TEPCO Fukushima dai-ichi nuclear power plant (FDNPP), leading to emission of huge amount of radionuclides to the environment. They have been transported and diffused by atmospheric motion, depositing them to soil and vegetation. Deposited radionuclides are dynamically shifted in the earth environment; atmosphere, soil, inland water, ocean, and ecosystem. To understand this dynamic shift in the environment and for the long-term prediction of the disaster by the radionuclides, investigation and discussion based on not only the earth sciences including ecology but also on the radiochemistry and other related sciences.

In this session, various efforts to understand the dynamic behavior of radionuclides emitted from FDNPP accident in the earth system as well as to predict their influences on the environment. It is expected that this session will offer a good opportunity to discuss radionuclides in the earth environment from wide aspect and to exchange information in various research fields.

## [MAG33-P02]Depth profile of $^{90}\text{Sr}$ concentration in soil ar Kawamata and Namie Town, Fukushima Prefecture

\*Anna SUZUKI<sup>1</sup>, Zijiang Zhang<sup>1</sup>, Kazuhiko Ninomiya<sup>1</sup>, Yoshiaki Yamaguchi<sup>2</sup>, Junko Takahashi<sup>3</sup>, Yuichi Onda<sup>3</sup>, Atsushi Shinohara<sup>1</sup> (1.Department of Chemistry, Graduate School of Science, Osaka University, 2.Radioisotope Research Center, Osaka University, 3.Center for Research in Isotopes and Environmental Dynamics, Tsukuba University)

Keywords:Sr-90, radioactive cesium, Fukushima Daiichi nuclear power plant, Depth Profile

Due to the accident of Fukushima Daiichi nuclear power plant (FDNPP) in Japan, large amounts of fission product in the nuclear reactor have been released into the atmosphere. Long lived radioactive nuclides such as  $^{90}\text{Sr}$  and radioactive cesium ( $^{134}\text{Cs}$  and  $^{137}\text{Cs}$ ), are the most significant radionuclides in the nuclear accident and the environmental fate of these nuclides have been attention. The activities of radioactive cesium can easily be determined from gamma-ray spectroscopy by Ge semiconductor detectors. Contrary to radioactive cesium, pure beta-ray emitting nuclide;  $^{90}\text{Sr}$  should be identified after chemical isolation. This is reason the studies on environmental behavior related to  $^{90}\text{Sr}$  were very limited compare to these of radioactive cesium. A part of the released radionuclides from the nuclear reactor deposits on the land area with rainfall. The environmental behavior of the radionuclides is different behaviors for downward movement due to the difference in on its chemical property<sup>[1]</sup>. The types of the soil and content of organic matter and clay minerals strongly affects the downward movement of radioactive nuclides. To estimate the long-term environmental behavior, it is necessary to evaluate the environmental dynamics of radionuclides in Fukushima prefecture. In this study, depth-profiling of  $^{90}\text{Sr}$  and radioactive cesium were investigated with different surrounding environments near the FDNPP. Soil samples were obtained at young aged cedar forests in Kawamata Town, and at the open land of Namie Town, Fukushima Prefecture, 2016. Because the radioactive nuclides were supplied from the organic phase to the soil with time passing, radioactivities in the litter was also identified for the sample in Kawamata town. For determination of radioactive cesium concentrations, gamma-ray measurement by high purity germanium

detectors were performed. For  $^{90}\text{Sr}$  identification, we adopted a multistage classical precipitation method to separate  $^{90}\text{Sr}$  fraction. To detect  $^{90}\text{Sr}$  activities, the growth of  $^{90}\text{Y}$ ; daughter nuclide of  $^{90}\text{Sr}$  was measured from Cherenkov light counting. The yield of  $^{90}\text{Sr}$  was determined from the amount of  $^{\text{nat}}\text{Sr}$  added before the chemical separation as a carrier by ICP-MS measurement.

[1] S. Forsberg et al., J. Environ. Radioactivity., 2000, 50, 235-252