

---

[EJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-AG Applied Geosciences

## [M-AG33] Dynamics of radionuclides emitted from Fukushima Dai-ichi Nuclear Power Plant in the environment

convener: Kazuyuki Kita (Faculty of Science, Ibaraki University), Yuichi Onda (Center for Research on Isotopes and Environmental Dynamics, University of Tsukuba), SHINOHARA ATSUSHI (大阪大学, 共同), Daisuke Tsumune (Central Research Institute of Electric Power Industry)

Sun. May 20, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

The Great East Japan Earthquake caused the severe accident in TEPCO Fukushima dai-ichi nuclear power plant (FDNPP), leading to emission of huge amount of radionuclides to the environment. They have been transported and diffused by atmospheric motion, depositing them to soil and vegetation. Deposited radionuclides are dynamically shifted in the earth environment; atmosphere, soil, inland water, ocean, and ecosystem. To understand this dynamic shift in the environment and for the long-term prediction of the disaster by the radionuclides, investigation and discussion based on not only the earth sciences including ecology but also on the radiochemistry and other related sciences.

In this session, various efforts to understand the dynamic behavior of radionuclides emitted from FDNPP accident in the earth system as well as to predict their influences on the environment. It is expected that this session will offer a good opportunity to discuss radionuclides in the earth environment from wide aspect and to exchange information in various research fields.

---

## [MAG33-P04] Evaluation of trapping of suspended Cs-137 in reservoir lakes in river systems in Fukushima

\*Keisuke Taniguchi<sup>1</sup>, Takayuki KURAMOTO<sup>1</sup>, Hirotsugu Arai<sup>1</sup>, Yuichi Onda<sup>2</sup>, Kazuya Yoshimura<sup>3</sup>, Hugh Smith<sup>4</sup>, William Blake<sup>5</sup>, Takayuki Sato<sup>1</sup>, Saori Onuma<sup>1</sup> (1. Centre for Environmental Creation Fukushima Prefecture, 2. University of Tsukuba, 3. Japan Atomic Energy Agency, 4. University of Liverpool, 5. Plymouth University)

Keywords: Radiocaesium, Reservoirs, TEPCO's Fukushima Daiichi Nuclear Power Plant accident

Radiocaesiums released by TEPCO's Fukushima Daiichi Nuclear Power Plant accident were deposited and adsorbed to fine particles of the surface soils such as silts and clays. The contaminated soils were eroded by rainfall events, and then transported through river systems. We have measured the activity concentrations and fluxes of suspended Cs-137 at 30 monitoring points in located in 10 river systems in the area affected by the accident. In order to evaluate the trapping effect of the large reservoir lakes on the riverine transport of suspended Cs-137, the suspended Cs-137 flux normalized by the water discharge and initial deposition of Cs-137 in the watershed ( $L/QD$ ) were compared with the coverages of the land use types and the proportion of the catchment area by reservoir lakes to the total basin area.

$L/QD$  values at the sites without reservoir(s) showed positive correlations with the coverage of paddy field, urban area and farmland, and a negative correlation with the forest coverage.  $L/QD$  values at the sites with reservoir(s) were 48 ~ 84 % of the values estimated by the land use coverages, and showed negative correlation with the proportion of the catchment area by reservoir lakes to the total basin area.