

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-GI General Geosciences, Information Geosciences & Simulations

[M-GI25]Environmental changes in mountainous area

convener:Keisuke Suzuki(Department of Environmental Sciences, Faculty of Science, Shinshu University), Yoshihiko Kariya(Department of Environmental Geography, Senshu University), Chiyuki Narama(新潟大学理学部理学科, 共同), Akihiko SASAKI(Department of Geography and Environmental Studies, Kokushikan University)
Tue. May 22, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)
Mountainous areas provide water resources to the populated downstream areas, protecting the diversity of ecosystem and providing tourism attraction. To access the mountain environment changes and mitigate the impacts of global warming influences, a cross-cutting session is proposed to share the scientific knowledge among various fields; such as climatology, hydrology, geography, glaciology, water/carbon/material cycle, eco-diversity, etc.

[MGI25-P08]Trench excavation in the lacustrine sediments in a dammed lake formed by the Dondokosawa rock avalanche and high-resolution landslide chronology using oxygen isotopic dendrochronology

*Ryuji Yamada¹, Yoshihiko Kariya², Kimio Inoue³, Takashi Kimura¹, Masaki Sano⁴, Zhen Li⁵, Akane Tsushima⁵, Takeshi Nakatsuka⁵ (1.National Research Institute for Earth Science and Disaster Resilience, 2.Senshu University, 3.Sabo Frontier Foundation, 4.Waseda University, 5.Research Institute for Humanity and Nature)
Keywords:Akaishi Mountains, rock avalanche, dammed lake, oxygen isotopic dendrochronology, high-resolution chronology

Dendrochronology, which has been mainly used in archaeology, is the science or technique of dating events, environmental changes in the past by using the characteristic patterns of annual growth rings in trees based on the fact that features of tree-rings are individually synchronized reflecting the annual changes in climatic factors such as temperature and precipitation. In recent years, applicable areas of dendrochronology have extensively been expanded by measuring instrumentally the oxygen isotopes of cellulose extraction instead of the widths as an index of changes. It is necessary to establish regional standard chronology (master chronology) as long as possible for dating of woods from various periods for both of widths and isotopes, and therefore the dendrochronological analysis can be performed in the latest ca. 3000 years under the status quo in Japan. Because climate sensitivity of tree-ring width depends on tree species, it is difficult to apply the master chronology made for a specific species to analysis for another species. On the other hand, tree-ring cellulose oxygen isotopic ratio reflects only two meteorological parameters (precipitation isotope ratios and relative humidity during growing season), and its variation shows high coincidence among different species. Oxygen isotopic dendrochronology consequently enables us to reconstruct palaeo climate annually (or seasonally if tree-rings sliced).

The Dondokosawa rock avalanche (DRA) blocked the main stream of the Dondokosawa River and formed dammed lakes besides the debris (Kariya, 2012). Many woody plant fossils were observed there. Yamada et al. (2018) applied the dendrochronological analysis using tree-ring oxygen isotope ratios to two tree trunks preserved in fairly good condition in the lacustrine sediments, and obtained their death years of around AD885 and AD888. Compared with historical records of large earthquakes in old documents such as *Shoku Nihongi* (a chronicle of Japan edited in AD 797), these ages implied that DRA occurred at the year of AD887 Ninna (Goki-Shichido) earthquake, which was one of the gigantic ocean-trench earthquakes along the Suruga

and Nankai Troughs off central Japan, or at a couple of years later than this earthquake. Because several active faults in the southern segments of Itoigawa-Shizuoka Tectonic Line (ISTL) active fault zone run through in this area, one of these fault could remotely be triggered by the Ninnu great interplate earthquake, like the case of the northern Nagano Prefecture earthquake (on March 12, 2011; $M = 6.7$) after the 2011 Tohoku earthquake (on March 11, 2011; $M = 9.0$).

We conducted an additional trench excavation in the lacustrine sediments in one of dammed lakes along the Dondokosawa River, and some buried tree trunk samples could be taken there in August 2016. Their conditions were not good because tree barks are detached while sampling. Dendrochronological dating yielded estimated years of AD830-880 for the outer most tree-rings remained (death years should exceed them) of the most of trunk samples, which were consistent with previous estimates. Outlines of the trench excavation and details of dendrochronological analysis are to be reported.

References: Kariya (2012) *Trans. Jpn. Geomorphol. Union* 33, 297-313.; Yamada et al. (2018) *Quat. Geochr.* 44, 47-54.