
[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS10]Paleoclimatology and paleoceanography

convener:Yusuke Okazaki(Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University), Atsuhiko Isobe(Research Institute for Applied Mechanics, Kyushu University), Akihisa Kitamura(静岡大学理学部地球科学教室, 共同), Masaki Sano(Faculty of Human Sciences, Waseda University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Past environmental changes and events at multi-decadal to tectonic timescale toward an understanding of Earth climate system by an integration of terrestrial and marine proxy studies and numerical modeling will be discussed. We welcome a variety of paleo-environmental studies from a wide range of background. In particular, a series of presentations relating to the Anthropocene will be planned. This is a merged session of A-OS31 "Linkage between oceanography and paleoceanography in marginal, shelf and coastal oceans" and M-IS23 "Paleoclimatology and paleoceanography" sessions at JPGU 2017. We hope that this session will provide an opportunity to promote communication between participants from multidisciplinary field.

[MIS10-P03]Reconstruction of sea-surface temperature change during the late Pliocene to early Pleistocene using Garda Drift sediment in the North Atlantic Ocean

Yuki Hikasa², *Toshiro Yamanaka¹, Hitoshi CHIBA², Tatsuya Hayashi³, Masao Ohno³ (1.Tokyo University of Marine Science and Technology, 2.Okayama University, 3.Kyushu University)

Keywords:Atlantic Meridional Overturning Circulation, Sea-surface temperature, Pliocene-Pleistocene boundary

In order to reveal the relationship between sea-surface temperature (SST) of high latitude in North Atlantic Ocean and current strength of North Atlantic Deep Water (NADW) associated with Atlantic Meridional Overturning Circulation (AMOC), which is considered responsible for 1/3 of global northward heat transport, a high-resolution reconstruction of SST has been performed using the core samples obtained at IODP Site U1314 in the Gardar Drift (56°21.9'N, 27°53.3'W in a water depth of 2820 m). We analyzed long-chain alkenones for reconstruction of SST in the subsamples of the age span from 2.74 to 2.42 Ma (late Pliocene to early Pleistocene). The mean sedimentation rate of the Gardar drift is considered very high (about 10 cm/kyr), therefore, the sediment obtained from there is expected to reconstruct paleoclimatic and paleoceanographic changes relevant to variation in strength of AMOC at a high-resolution.

As a result, the amplitude of SST variations were enhanced after the boundary between Pliocene and Pleistocene, and the spike-like falls of SST were detected at 2.62Ma, 2.52Ma, and 2.43Ma, corresponding to MIS104, MIS100, and MIS 96, respectively. Compared the SST variations with the numbers of ice rafted debris (IRD) and variations in strength of deep water current (those data will be presented by Hayashi et al. in this meeting) and global ice volume (LR04), the amplitudes were different each other, but the timings of changes were in agreement with each other. On the other hand, some changes appeared in both SST and the strength of deep water current were not appeared in LR04, suggesting that SST variation observed at U1314 Site was influenced by the variation in strength of NADW, that is the strength of AMOC.