

---

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

## [M-IS10]Paleoclimatology and paleoceanography

convener:Yusuke Okazaki(Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University), Atsuhiko Isobe(Research Institute for Applied Mechanics, Kyushu University), Akihisa Kitamura(静岡大学理学部地球科学教室, 共同), Masaki Sano(Faculty of Human Sciences, Waseda University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Past environmental changes and events at multi-decadal to tectonic timescale toward an understanding of Earth climate system by an integration of terrestrial and marine proxy studies and numerical modeling will be discussed. We welcome a variety of paleo-environmental studies from a wide range of background. In particular, a series of presentations relating to the Anthropocene will be planned. This is a merged session of A-OS31 "Linkage between oceanography and paleoceanography in marginal, shelf and coastal oceans" and M-IS23 "Paleoclimatology and paleoceanography" sessions at JPGU 2017. We hope that this session will provide an opportunity to promote communication between participants from multidisciplinary field.

---

## [MIS10-P06]Biomarker analysis of the uncorrelated Early-Pleistocene marine clay layer of the Osaka Group in the Kishiwada City, Osaka Prefecture

\*Hideto Nakamura<sup>1</sup>, Takuto Ando<sup>2</sup> (1.Faculty of Science, Osaka City University, 2.Arctic Research Center, Hokkaido University)

Keywords:Long chain diols, n-alkanes, Osaka Group, Sedimentary environment, Marine clay

The Osaka Group is characterized by thick sedimentation of Plio-Pleistocene sediments (>2000 m in the basin center) due to rapid subsidence. The lowest part of the Osaka Group is freshwater deposits consist of sand, silt and mud, which frequently intercalated with gravel layer. In the lower to the upper part of the Osaka Group, marine clay layers repeatedly deposit spanning wide area of the Osaka Basin, and has been given symbols as Ma-1, Ma0, Ma1...up to Ma-13. Deposition of marine clay layers are affected by eustatic sea-level changes, and the Ma-1 is considered to be the earliest significant marine incursion through the Kii Channel into a large area of Osaka Basin. Recently, fresh outcrop including unidentified marine-like clay layer (MLC) was newly exposed near the Tonboike Park, Kishiwada City. This MCL is presumed to be older than Ma1, however, have not been correlated to the certain known marine clay layer due to a lack of key bed within the outcrop.

In order to examine the source of organic matter and to reconstruct the changes in depositional environment during the possible sea-level rise event, sediments including MCL layer were collected from the outcrop and analyzed for biomarker compositions. Composition of long-chain diols in the upper part of the MCL layer was characterized by a predominance of C<sub>30</sub> diols over C<sub>32</sub> diols, which correspond to the central to inner part of the recent Osaka Bay. In contrast, the lower part of MCL was rich in C<sub>32</sub> diols indicating significant influence of freshwater input. Underlying fluviolimnic sediments were characterized with the predominance of C<sub>32</sub> 1,15-diol indicating freshwater environment. Steroid compositions (the relative amount of C<sub>27</sub>, C<sub>28</sub>, C<sub>29</sub> stanols) indicated the varying input of terrestrial to aquatic organic matter, which is concordant with both the results of long-chain diols and the lithofacies observation. The *n*-alkane index TAR reconstructed variation in the contribution of aquatic OM along with continual input of terrestrial plant wax. In addition, elevated *Paq* value was noted at the lower part of the MCL. The combination of multiple biomarker indices suggests that (1) the upper part of the marine like clay layer is deposited under marine environment, and (2) potential development of the blackish aquatic environment with high input of aquatic macrophytes at the earliest

stage of the history of Osaka Bay marine environment before the typical marine mud layer deposition.