

[JJ] Evening Poster | M (Multidisciplinary and Interdisciplinary) | M-IS Intersection

[M-IS10]Paleoclimatology and paleoceanography

convener:Yusuke Okazaki(Department of Earth and Planetary Sciences, Graduate School of Science, Kyushu University), Atsuhiko Isobe(Research Institute for Applied Mechanics, Kyushu University), Akihisa Kitamura(静岡大学理学部地球科学教室, 共同), Masaki Sano(Faculty of Human Sciences, Waseda University)

Wed. May 23, 2018 5:15 PM - 6:30 PM Poster Hall (International Exhibition Hall7, Makuhari Messe)

Past environmental changes and events at multi-decadal to tectonic timescale toward an understanding of Earth climate system by an integration of terrestrial and marine proxy studies and numerical modeling will be discussed. We welcome a variety of paleo-environmental studies from a wide range of background. In particular, a series of presentations relating to the Anthropocene will be planned. This is a merged session of A-OS31 "Linkage between oceanography and paleoceanography in marginal, shelf and coastal oceans" and M-IS23 "Paleoclimatology and paleoceanography" sessions at JPGU 2017. We hope that this session will provide an opportunity to promote communication between participants from multidisciplinary field.

[MIS10-P10]Environmental fluctuations recorded in dark-light alternations of sediment in the southern Japan Sea offshore Wakasa Bay

*Yusuke Narita¹, Masahiro Yokoyama¹, Takuya Sagawa², Akiko S. Goto², Takashi Hasegawa², Yoshimi Kubota³ (1.Kanazawa University, College of Science and Engineering, 2.Kanazawa University, Institute of Science and Engineering, 3.National Museum of Nature and Science)

Keywords:Last glacial period, Japan Sea, Organic carbon, East Asian summer monsoon, Dansgaard-Oeschger cycle

Alternations of dark and light color are observed in the last glacial sediments of the Japan Sea, and change in brightness of sediments resembles millennial cycle of $\delta^{18}O$ in the Greenland ice core (Dansgaard-Oeschger cycle: DO cycle). Although brightness (L^*) of Japan Sea sediment mainly reflects total organic carbon (TOC) content, as TOC content changes due to various factors, such as primary production in the ocean surface and post-depositional degradation etc., it is difficult to reconstruct marine environmental change only by TOC content in sediments. Chemical composition of planktonic foraminiferal tests record the information of surface water at the time of shell formation, and the oxygen isotope ratio and Mg/Ca are excellent proxies for reconstructing past ocean temperature. Therefore, paired analysis of TOC and planktonic foraminiferal geochemistry would give an insight into surface process and its relation to TOC variation in the Japan Sea sediments. However, there have been no attempt to reconstruct millennial-scale surface water temperature change corresponding to the TOC content variation in the Japan Sea. In this study, we aimed to understand the change of the marine environment of Japan Sea in conjunction with the DO cycle by reconstructing the surface water temperature and analyzing TOC content at high resolution in a piston core sediment (KR-15-10-WB-6_PC, water depth 845 m) collected off the Wakasa Bay in the southern Japan Sea.

Variations of L^* and TOC contents of WB6_PC are highly correlated as previously reported by several studies. Changes in sea surface temperature (SST) inferred from oxygen isotope and Mg/Ca of planktonic foraminifera also show good correlation with TOC, and warmer temperature corresponds to higher TOC. Therefore, it is suggested that the change in TOC content is closely related to surface water property change, implying primary production co-varied with SST. Coupling of TOC and SST could be due to the millennial-scale variation of the East Asian summer monsoon (EASM). The strength of EASM could change nutrient influx into Japan Sea via terrestrial nutrient discharge from Changjiang River. Enhanced nutrient availability during

strong EASM periods would stimulate primary production. In addition, northward shift of the westerly jet axis during the strong EASM could affect SST in the Japan Sea due to expansion of subtropical warm air mass. Therefore, the EASM could be an important component linking climate of the Greenland and Japan Sea.